


## FRONT LABEL



<b>CONTROLS</b>	<input type="checkbox"/> NEW <input checked="" type="checkbox"/> REVISIONS <input type="checkbox"/> OTHER			approval sign-off: _____ date: _____	
	client: <b>IMAGE</b>			colors: <input checked="" type="checkbox"/> CMYK <input checked="" type="checkbox"/> SPOT COLORS	
	date submitted: <b>XX/XX/20</b>	version date: <b>03/15/21</b>	requested finish: <b>XX/XX/20</b>	PMS # 5425 <input checked="" type="checkbox"/>	PMS # 341 <input checked="" type="checkbox"/>
	file name: <b>IM125 - Image 100523494 All in One Lawn Weed Killer RTS 24oz</b>			PMS # 000 <input checked="" type="checkbox"/>	PMS # 000 <input checked="" type="checkbox"/>
	requester: <b>J. DIETZ</b>	designer: <b>SHAK</b>	version #: <b>18</b>	This color print is for content and position only. Use Contract Proof for Process-Color matching. Use Ink Drawdowns for Spot-Color matching.	

**CENTRAL**  
Garden & Pet

CENTRAL GARDEN & PET | V2.0 | 4/213

## BASE LABEL (BACK)



### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

##### CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants, shoes, socks and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

##### ENVIRONMENTAL HAZARDS

This product is toxic to marine/estuarine invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

This product may impact surface water quality due to runoff of rain water or spray drift. Under certain conditions this product may have a high potential for runoff into surface water for several months after application. This is especially true for poorly draining soils and soils with shallow ground water.

The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of labeled use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

#### FIRST AID

<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call poison control center or doctor immediately for treatment advice.</li><li>• Have a person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

#### HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information.

 **CENTRAL**  
GARDEN & PET  
1-800-265-0761  
gardendivision@central.com

Manufactured for:  
Central Garden & Pet, Garden Division  
1000 Parkwood Circle, Suite 700  
Atlanta, GA 30339

EPA Reg No. 279-3463-90098  
EPA Est. No. 59144-GA-1, 67572-GA-1, 67572-GA-2,  
74750-GA-1, 70815-GA-1, 5905-GA-1  
Subscript used is first letter of Run Code on container. © 2015, 2017, 2020, 2021  
Central Garden & Pet Company  
300523711 - 03/21 | IM25\_V18



## BOOK COVER

**STARTS  
WORKING  
IMMEDIATELY**



### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.


#### APPLICATION INSTRUCTIONS AND PRECAUTIONS

<b>WHERE TO USE</b> 	<b>Lawns:</b> <b>Well established turfgrasses.</b> <b>Cool Season Grasses:</b> Fine & Tall Fescue, Perennial Ryegrass, Kentucky & Rough Bluegrass. <b>Warm Season Grasses:</b> Common & Hybrid Bermudagrass, Buffalograss, Centipedegrass, Seashore Paspalum, Zoysiagrass. <b>Do not use on St. Augustinegrass.</b>
<b>KILLS</b> 	Sedges, Grasses and Broadleaf Weeds: Yellow Nutsedge, Kyllinga, Crabgrass, Foxtail, Ground Ivy, Wild Violet, Plantain, Dandelion, Clover, Dollarweed, Spurge, Woodsorrel, Knotweed, Chickweed, Curly Dock, Wild Onion, Wild Garlic, and other listed weeds.
<b>WHEN TO USE</b>	When weeds are small and actively growing. For best performance, apply when temperatures are between 55°F - 85°F.
<b>HOW MUCH TO USE</b>	4 fl oz per 1 gallon of water per 1,000 sq.ft. Spot spray weed directly until thoroughly covered. Some weeds may require repeat application. (See detailed instructions.)
<b>COVERAGE</b>	24 fl.oz. covers up to 6,000 sq.ft.

See inside booklet for additional Precautionary Information, First Aid and complete Directions For Use.



**Questions & Comments: Call 1-800-265-0761 or visit our website at [www.imageforweeds.com](http://www.imageforweeds.com)**

300523711 - 3/21 | IM125\_V18  
  
8 13576 00690 7

Manufactured for:  
Central Garden & Pet, Garden Division  
1000 Parkwood Circle, Suite 700  
Atlanta, GA 30339  
1-800-265-0761  
[gardendivision@central.com](mailto:gardendivision@central.com)

EPA Reg No. 279-3463-90098  
EPA Est. No. 59144-GA-1, 67572-GA-1, 67572-GA-2,  
74750-GA-1, 70815-GA-1, 5905-GA-1  
Subscript used is first letter of Run Code on container.  
Image, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

# **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

## **PRODUCT INFORMATION**

This product selectively controls tough weeds like crabgrass, chickweed, dandelion, clover, ground ivy, purple<sup>†</sup> and yellow nutsedge (nutgrass), Kyllinga, wild onion and garlic, broadleaf plantain, purslane, wild violet, spurge, and other listed annual, biennial, and broadleaf weeds in a variety of lawns, including:

<b>Cool Season Grasses</b>	<b>Warm Season Grasses</b>
Tall & Fine Fescue, Kentucky Bluegrass, Rough Bluegrass, Ryegrass (perennial)	Common & Hybrid Bermudagrass, Zoysiagrass, Centipedegrass, Buffalograss, Seashore Paspalum

<sup>†</sup>Not for use in California

This product has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties have been evaluated. Some turf varieties are more susceptible to temporary turf discoloration than others. It is recommended to treat a small area prior to treatment of larger areas.

This unique formula causes visual symptoms to develop quickly within 3 to 7 days after application. This product involves uptake by both weed roots and shoots. Complete kill should occur in 2-3 weeks. This product also controls yellow nutsedge tubers (nutlets) in the soil over time.

A postemergence application of this product is improved when adequate soil moisture is present at application.

## **HOW TO APPLY:**

### **Ready to Spray Hose-End Sprayer**

1. Measure the entire area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet.
2. See below for directions for the ready-to-use hose-end sprayer.
3. Spray evenly over the entire measured area.

<b>Amount of Product</b>	<b>Provides Coverage of Area Size:</b>
1 fl oz	250 sq.ft.
2 fl oz	500 sq.ft.
4 fl oz	1000 sq.ft.
6 fl oz	1500 sq.ft.
8 fl oz	2000 sq.ft.
24 fl oz	6000 sq.ft.

## Hose End Sprayer Connection

1. Shake well.
2. Make sure control lever is in the OFF position. Attach to hose.
3. Turn on water.

## Spray

1. Aim towards direction of application and away from you.
2. Turn control lever to ON to begin spraying.
3. Spray evenly and as directed for specified type of treatment area.

## Spray

1. When finished turn control lever to OFF.
2. Turn faucet OFF. Discharge residual water pressure by turning control lever ON and OFF again.
3. Disconnect Sprayer and Store / Discard as instructed.

**COVERAGE:** 4 fl.oz. per 1000 sq. ft.

## APPLICATION INSTRUCTIONS:

- Best weed control results will be obtained when no rainfall or irrigation occurs within 24 hours after application.
- Listed sedges and grassy weeds: Spray directly on emerged weeds. This product is effective against newly emerged and established sedges and grassy weeds.
- Listed broadleaf weeds: Spray directly on newly emerged weeds.
- For best results, apply to weeds when less than 3 inches high.
- Apply in calm weather.
- Begin spraying at the farthest point of the treatment area and work backward to avoid contact with wet treated surfaces.
- Good spray coverage is needed for optimum control.
- Avoid overlap spraying.
- Avoid side to side or swinging arm motion, as it can result in uneven coverage.
- If listed weeds are not controlled by the initial application, apply a second application 4 to 6 weeks after the initial treatment.
- Water lawn with at least 0.5 inches within 7 days (but not sooner than 1 day after application) if no rainfall or irrigation occurs within 7 days after application.

## APPLICATION RESTRICTIONS

Keep children and pets out of the treated area until sprays have dried.

Do not spray over the top of flowers, ornamentals (including herbaceous) or garden vegetables.

Do not use on turfgrasses other than those listed on the label.

Do not use in greenhouses.

Do not graze or feed animals with treated grass clippings.

Do not apply to newly planted, plugged, or sodded turfgrass.

## APPLICATION PRECAUTIONS

Use only on well established non-stressed turfgrass. Applications to stressed plants (turfgrass or target weeds) may result in temporary discoloration or poor weed control. Allow turfgrass to develop good root system before applying.

If discoloration occurs, treated turfgrass will recover with new growth and discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turfgrass growth.

Avoid applying when windy to prevent product drifting to other plants.

Avoid reseeding, winter overseeding, or plugging lawns for at least 4 weeks after treatment.

### Grass and sedge weed species controlled or suppressed by this product:

Common Name	Scientific Name	Common Name	Scientific Name
Barneygrass	<i>(Echinochloa crus-galli)</i>	Kyllinga, fragrant	<i>(Kyllinga sesquiflorus)</i>
Broadleaf Signalgrass	<i>(Brachiaria platyphylla)</i>	Kyllinga, tufted	<i>(Kyllinga pumila)</i>
Crabgrass, Large	<i>(Digitaria sanguinalis)</i>	Japanese stiltgrass	<i>(Microstegium vimineum)</i>
Crabgrass, Smooth	<i>(Digitaria ischaemum)</i>	Junglerice	<i>(Echinochloa colona)</i>
Deergrass	<i>(Panicum cladestinum)</i>	Nutsedge, purple <sup>†</sup>	<i>(Cyperus rotundus)</i>
Goosegrass	<i>(Eleusine indica)</i>	Nutsedge, yellow	<i>(Cyperus esculentus)</i>
Foxtail, giant	<i>(Setaria faberi)</i>	Sedge, annual	<i>(Cyperus compressus)</i>
Foxtail, green	<i>(Setaria viridis)</i>	Sedge, globe	<i>(Cyperus globulosus)</i>
Foxtail, yellow	<i>(Setaria glauca)</i>	Sedge, cylindrical	<i>(Cyperus retrorsus)</i>
Kyllinga, cocks-comb	<i>(Kyllinga squamulata)</i>	Sedge, Surinam	<i>(Cyperus surinamensis)</i>
Kyllinga, green	<i>(Kyllinga brevifolia)</i>	Sedge, Texas	<i>(Cyperus polystachyos)</i>
Kyllinga, false green	<i>(Kyllinga gracillima)</i>		

For optimal control of Goosegrass, Junglerice, or Purple Nutsedge<sup>†</sup>, repeat treatment every 3-4 weeks.

<sup>†</sup>Not for use in California

### Broadleaf weeds controlled or suppressed by this product:

Common Name	Scientific Name	Common Name	Scientific Name
Bedstraw, catchweed	<i>(Galium aparine)</i>	Carpetweed	<i>(Mollugo verticillata)</i>
Beggarticks <sup>†</sup>	<i>(Bidens frondosa)</i>	Chickweed, common	<i>(Stellaria media)</i>
Beggarweed, Florida	<i>(Desmodium tortuosum)</i>	Chickweed, mouseear	<i>(Cerastium vulgatum)</i>
Bindweed, field	<i>(Convolvulus arvensis)</i>	Cinquefoil	<i>(Potentilla spp.)</i>
Bittercress	<i>(Cardamine spp.)</i>	Clover	<i>(Trifolium spp.)</i>
Black medic	<i>(Medicago lupulina)</i>	Clover, alyce	<i>(Alysicarpus ovalifolius)</i>
Buttercups	<i>(Ranunculus spp.)</i>	Clover, hop	<i>(Trifolium campestre)</i>
Buttonweed, common	<i>(Diodia teres)</i>	Clover, red	<i>(Trifolium pretense)</i>
(poorjoe)		Clover, white	<i>(Trifolium repens)</i>
Carolina geranium	<i>(Geranium carolinianum)</i>	Cudweed	<i>(Gnaphalium spp.)</i>

**Broadleaf weeds controlled or suppressed by this product (cont'd):**

Common Name	Scientific Name	Common Name	Scientific Name
Daisy, English	( <i>Bellis perennis</i> )	Pigweed, Smooth	( <i>Amaranthus hybridus</i> )
Dandelion	( <i>Taraxacum officinale</i> )	Pigweed, Tumble	( <i>Amaranthus albus</i> )
Dandelion, false	( <i>Microseris cuspidate</i> )	Pineapple weed	( <i>Matricaria matricariodes</i> )
Dayflower, common	( <i>Commelina diffusa</i> )	Plantain, Broadleaf	( <i>Plantago comanalata</i> )
Dock, Curly	( <i>Rumex crispus</i> )	Plantain, Buckhorn	( <i>Plantago lanceolata</i> )
Dollarweed	( <i>Hydrocotyle spp.</i> )	Plantain, Narrowleaf	( <i>Plantago narrowifolia</i> )
(pennywort)		Puncturevine	( <i>Tribulus terrestris</i> )
Eclipta	( <i>Eclipta prostrata</i> )	Purslane, common	( <i>Portulaca oleracea</i> )
Evening primrose	( <i>Oenothera biennis</i> )	Pusley, Florida	( <i>Richardia scabra</i> )
Fiddleneck	( <i>Amsinckia spp.</i> )	Redweed	( <i>Melochia corymbifolia</i> )
Filaree	( <i>Erodium spp.</i> )	Shepherd's-Purse	( <i>Capsella bursa-pastoris</i> )
Filaree, redstem	( <i>Erodium cicutarium</i> )	Smartweed,	( <i>Polygonum</i>
Gallinsoga	( <i>Gallinsoga ciliata</i> )	Pennsylvania	<i>pennsylvanicum</i> )
Goldenrod	( <i>Solidago spp.</i> )	Sorrel, Red	( <i>Rumex acetosella</i> )
Ground ivy	( <i>Glechmea hederacea</i> )	Speedwell	( <i>Veronica spp.</i> )
Groundsel, common	( <i>Senecio vulgaris</i> )	Speedwell, slender	( <i>Veronica filiformis</i> )
Henbit	( <i>Lamium amplexicaule</i> )	Speedwell, thymeleaf	( <i>Veronica serpyllifolia</i> )
Knapweed	( <i>Scleranthus annuus</i> )	Spurge, (annuals)	( <i>Euphorbia spp.</i> )
Knotweed, prostrate	( <i>Polygonum aviculare</i> )	Spurge, prostrate	( <i>Euphorbia humifusa</i> )
Kochia	( <i>Kochia scoparia</i> )	Spurge, spotted	( <i>Euphorbia maculata</i> )
Lambsquarters,	( <i>Chenopodium album</i> )	Star-of-Bethlehem	( <i>Ornithogalum</i>
common			<i>umbellatum</i> )
Lawn burweed	( <i>Soliva pterosperma</i> )	Velvetleaf	( <i>Abutilon theophrasti</i> )
(Spurweed)		Violet, wild	( <i>Viola pratensis</i> )
Lespedeza, common	( <i>Lespedeza striata</i> )	Violet,	( <i>Viola rafinesquii</i> )
London Rocket	( <i>Sisymbrium irio</i> )	Johnny-Jump-Up	
Mallow, common	( <i>Malva neglecta</i> )	Wild garlic	( <i>Allium vineale</i> )
Morning glory	( <i>Ipomea spp.</i> )	Wild onion	( <i>Allium canadense</i> )
Morning glory,	( <i>Ipomoea hederacea</i> var.	Woodsorrel, creeping	( <i>Oxalis corniculata</i> )
entireleaf	<i>integrifolia</i> )	Woodsorrel, yellow	( <i>Oxalis stricta</i> )
Morning glory, ivyleaf	( <i>Ipomoea hederacea</i> )		
Morning glory,	( <i>Ipomoea wrightii</i> )		
palmleaf			
Morning glory, pitted	( <i>Ipomoea lacunosa</i> )		
Morning glory, tall	( <i>Ipomoea purpurea</i> )		
Parsley piert	( <i>Alchemilla arvensis</i> )		
Pigweed, Redroot	( <i>Amaranthus retroflexus</i> )		

\*Not for use in California

Certain listed weeds may require more than one application to achieve season-long control.

# STORAGE AND DISPOSAL

## Pesticide Storage

Keep from freezing. Do not store below 32°F. Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat.

**In case of a spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Central Garden & Pet, Garden Division: 1-800-265-0761.**

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

## Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

## Container Handling

Non-refillable container. Do not reuse or refill this container. If empty: Offer for recycling, if available, or dispose of in trash.

If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NOTICE:** To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.



## NOTES

---

---

---

---

---

---

---

---

## NOTES

---

---

---

---

---

---

---

---