

Group 3 11 Fungicide



PRO Granular Fungicide

For disease control and plant health in turfgrass

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

See attached booklet for complete **First Aid, Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty.**

In case of emergency endangering life or property involving this product, call day or night 1 (888) 270-3714.

FIRST AID:

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. **DO NOT** give anything by mouth to an unconscious person.

If in eyes: Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eyes. Call a poison control center for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact for emergency medical treatment information: 1-877-465-5161.

Precautionary Statements:

Hazards to Humans and Domestic Animals:

CAUTION. Harmful if absorbed through skin. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, pants, socks, shoes, gloves and protective eyewear.

Environmental Hazards: DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

Directions For Use: It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. Observe all precautions and limitations in this label and the labels of products used in combination with **Scotts® Pro Granular Fungicide**. The use of **Scotts® Pro Granular Fungicide** not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions and Conditions of Sale and Warranty are to be followed.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited. **Pesticide Storage: DO NOT** store above 120° F. Store in original containers and keep closed. Store in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility. **CONTAINER HANDLING: Nonrefillable Container. DO NOT** reuse or refill this container. After completely emptying container into application equipment, dispose of empty bag in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

37612-1 LB17293

The Scotts Company
14111 Scottslawn Road
Marysville, OH 43041

Net Content: 12 pounds



Group 3 11 Fungicide

Scotts **PRO**
Granular Fungicide

For disease control and plant health in turfgrass

Active Ingredients:

Pyraclostrobin0.38%

Triticonazole0.43%

Other Ingredients:99.19%

Total:100.00%

0.0038 lb of pyraclostrobin in 1 lb of **Scotts Pro Granular Fungicide**

0.0043 lb of triticonazole in 1 lb of **Scotts Pro Granular Fungicide**

EPA Reg. No. 538-345

EPA Est. No. 51036-GA-001

KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

See inside for complete **First Aid, Precautionary Statements, Directions For Use and Conditions of Sale and Warranty**

In case of emergency endangering life or property involving this product, call day or night
1-888-270-3714

Net Contents: 12 pounds

The Scotts Company
14111 Scottslawn Road
Marysville, OH 43041

37612-1 LB17293



FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eyes. • Call a poison control center for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-465-5161 for emergency medical treatment information.	

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Harmful if absorbed through skin. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, pants, socks, shoes, gloves and protective eyewear.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Observe all precautions and limitations in this label and the labels of products used in combination with **Scotts® Pro Granular Fungicide**. The use of **Scotts® Pro Granular Fungicide** not consistent with this label can result in injury to crops, animals, or persons.

Keep containers closed to avoid spills and contamination.

Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** are to be followed.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves (made of any waterproof materials)
- Shoes plus socks
- Protective eyewear

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. **DO NOT** enter or allow others to enter the treated area until dusts have settled.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

DO NOT store above 120° F. Store in original containers and keep closed. Store in a cool, dry place.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

Container Handling

Nonrefillable Container. **DO NOT** reuse or refill this container. After completely emptying container into application equipment, dispose of empty bag in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

In Case of Emergency

- In case of large-scale spillage regarding this product, call:
- Chemtrec 1-800-424-9300

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- The Scotts Company 1-877-465-5161

Steps to be taken in case this material is released or spilled:

- Dike and contain spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing, and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep spill out of all sewers and open bodies of water.

Product Information

Apply **Scotts® Pro Granular Fungicide** to turfgrass as a broad-spectrum preventive fungicide for the control of many important turfgrass diseases. Preventive applications optimize disease control and may result in improved plant health. **Scotts® Pro Granular Fungicide** contains pyraclostrobin, the active ingredient in **Insignia® fungicide**, and triticonazole, the active ingredient in **Trinity® fungicide**. This combination product, with two modes of action, is an excellent resistance management tool.

Application Information

Scotts® Pro Granular Fungicide is a broad-spectrum fungicide for the control of many important diseases of turfgrass. Apply **Scotts® Pro Granular Fungicide** with spreader equipment commonly used for ground application. Proper calibration of equipment to give good coverage is essential for disease control. Apply **Scotts® Pro Granular Fungicide** preventively. **DO NOT** exceed the specified application rate or fail to comply with use restrictions listed in the **Restrictions and Limitations** section.

Use Sites

Turfgrass

Use **Scotts® Pro Granular Fungicide** for disease control in the following turf use sites:

- golf courses
- institutional, commercial and residential lawns
- sod farms
- parks
- recreational areas including sports fields
- cemeteries

Tolerant Turfgrass Species

- | | |
|------------------------------------|-----------------------|
| • Bentgrass, colonial | • Dichondra |
| • Bentgrass, creeping | • Fescue, fine |
| • Bermudagrass, common | • Fescue, tall |
| • Bermudagrass, hybrid | • Paspalum, seashore |
| • Bluegrass, annual (Poa annua) | • Ryegrass, annual |
| • Bluegrass, Kentucky | • Ryegrass, perennial |
| • Bluegrass, rough (Poa trivialis) | • St. Augustinegrass |
| • Buffalograss | • Zoysiagrass |
| • Centipedegrass | |

Application Information *(continued)*

Due to variability within turfgrass species and application techniques, neither the manufacturer nor the seller has determined whether **Scotts® Pro Granular Fungicide** can safely be used on all turf grasses under all conditions. The user must determine if **Scotts® Pro Granular Fungicide** can be used safely before broadscale use. Apply the specified use rate of **Scotts® Pro Granular Fungicide** on a small test area under conditions expected to be encountered. Monitor for any adverse effects during a 14-day period after application. **DO NOT** use on ultradwarf bermudagrass varieties.

When used according to label instructions, Scotts® Pro Granular Fungicide is effective in controlling the following diseases

- | | |
|-----------------------------------|-----------------------------------|
| • Anthracnose | • Pink patch |
| • Bentgrass dead spot | • Pink snow mold |
| • Bermudagrass decline | • Powdery mildew |
| • Brown patch | • Pythium blight |
| • Brown ring patch (Waitea patch) | • Pythium root dysfunction |
| • Dollar spot | • Rapid blight |
| • Fairy ring | • Red thread |
| • Fusarium patch | • Rhizoctonia leaf or sheath spot |
| • Gray leaf spot | • Rust |
| • Gray snow mold | • Summer patch |
| • Large patch | • Take-all patch |
| • Leaf spot | • Yellow tuft (Downy mildew) |
| • Melting out | • Zoysia patch |
| • Necrotic ring spot | |

Resistance Management

The active ingredients in **Scotts® Pro Granular Fungicide** are pyraclostrobin, a **Group 11 Quinone Outside Inhibitor (QoI)** and triticonazole, a **Group 3 demethylation inhibitor (DMI)**. Fungal isolates resistant to **Group 11** (strobilurin or QoI) fungicides such as pyraclostrobin, azoxystrobin, trifloxystrobin and kresoxim-methyl, and **Group 3** (DMI) fungicides such as fenarimol, myclobutanil, triadimefon, propiconazole, tebuconazole, or metconazole may allow less sensitive strains of target fungi to build over time. This may result in reduced disease control. To limit the potential for the development of resistance alternate **Scotts® Pro Granular Fungicide** with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed. **DO NOT** make more than 2 sequential applications of **Scotts Pro Granular Fungicide**. **DO NOT** alternate **Scotts® Pro Granular Fungicide** with other **Group 11** fungicides.

Restrictions and Limitations

- **Maximum seasonal use rate** - **DO NOT** apply more than a total of 15 lbs/1000 sq ft of **Scotts® Pro Granular Fungicide** per year.
- Refer to the **Scotts® Pro Granular Fungicide Application Directions on Turfgrass** section for sequential application intervals for **Scotts® Pro Granular Fungicide**.
- **DO NOT** use on crops intended for food or feed use.
- **DO NOT** use this product to formulate or reformulate any other pesticide product.

Scotts® Pro Granular Fungicide Application Directions on Turfgrass

Apply **Scotts® Pro Granular Fungicide** at a use rate of 3.0 lbs product/1000 sq ft (131 lbs product/acre) on a 14-day to 28-day interval for the following diseases. **DO NOT** make more than 5 applications per year at the use rate of 3.0 lbs product/1000 sq ft (131 lbs product/acre). Applications made to control root and soilborne diseases such as fairy ring, Pythium root dysfunction, summer patch and take-all patch will require additional irrigation to move product into the soil.

Disease Pathogen	Application Interval (days)	Comments
Anthracnose <i>Colletotrichum graminicola</i>	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
Bentgrass dead spot <i>Ophiosphaerella agrostis</i>	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
Bermudagrass decline <i>Gaeumannomyces graminis</i> var. <i>graminis</i>	28	Aids in control of bermudagrass decline when integrated with appropriate cultural practices such as raised mowing height, proper fertilization, and core aeration. Make 1 application in the spring following greenup and a second application in the fall when air temperatures remain above 80° F and humidity is 75% or higher.
Brown patch <i>Rhizoctonia solani</i>	14 to 28	Apply when conditions are favorable for disease development.
Brown ring patch <i>Rhizoctonia circinata</i> var. <i>circinata</i> aka Waitea patch	14 to 28	Apply when early yellow ring development is symptomatic. Provide short irrigation cycle directly following treatment to move fungicide through thatch.
Dollar spot <i>Sclerotinia homoeocarpa</i>	14 to 28	Apply when conditions are favorable for disease development. Ring symptoms can take 2 to 3 weeks to disappear following application. For optimum results, apply a soil wetting agent to the intended treatment area prior to Scotts® Pro Granular Fungicide application.
Fairy ring various <i>Basidiomycete</i> fungi	14 to 28	Apply as soon as possible after fairy ring symptom development. Fairy ring symptoms can take 2 to 3 weeks to disappear following application. For optimum results, apply a soil wetting agent to the intended treatment area prior to Scotts® Pro Granular Fungicide application.
Fusarium patch (Pink snow mold) <i>Microdochium nivale</i>	28	Make 1 application in late fall just prior to snow cover.
Gray leaf spot <i>Pyricularia grisea</i>	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
Gray snow mold/Typhula blight <i>Typhula</i> spp.	28	Make 1 application in late fall just prior to snow cover.
Large patch <i>Rhizoctonia solani</i> AG2-2LP	14 to 28	Make 1 to 2 applications in the fall when conditions are favorable for disease development, repeat in spring prior to and following greenup.
Leaf spot <i>Bipolaris</i> spp., <i>Dreschlera</i> spp., <i>Exserohilum</i> spp.	14 to 28	Apply when conditions are favorable for disease development.
Melting out <i>Drechslera poae</i>	14 to 28	Apply when conditions are favorable for disease development.
Necrotic ring spot <i>Leptosphaeria korrae</i>	28	Apply when soil temperatures reach 60° F in early spring to summer and continue at 28-day intervals under conditions favorable for disease development. Provide additional irrigation following application.

(continued)

Scotts® Pro Granular Fungicide Application Directions on Turfgrass *(continued)*

Disease Pathogen	Application Interval (days)	Comments
Pink patch <i>Limonomyces roseipellis</i>	14 to 28	Apply when conditions are favorable for disease development, when night temperatures are 60° F to 70° F, and in periods of higher rainfall.
Pink snow mold (Fusarium patch) <i>Microdochium nivale</i>	28	Make 1 application in late fall just prior to snow cover.
Powdery mildew <i>Blumeria graminis</i>	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
Pythium blight <i>Pythium aphanidermatum</i> , <i>Pythium</i> spp.	14	Use preventively. Begin applications when conditions are favorable for fungal infection, normally under prolonged moist and humid conditions. Apply prior to disease symptom development for best control.
Pythium root dysfunction <i>Pythium volutum</i> spp., <i>Pythium</i> spp.	14	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development. DO NOT apply more than two strobilurin based fungicides sequentially prior to rotation to a different mode of action.
Rapid blight <i>Labyrinthula terrestris</i>	14 to 28	Apply when conditions are favorable for disease development which occurs normally on salinity stressed grasses. Occurs most commonly on cool season grasses used under overseeded conditions. Begin applications in fall and follow with spring applications as needed. DO NOT apply more than two strobilurin based fungicides sequentially prior to rotation to a different mode of action.
Red thread <i>Laetisaria fuciformis</i>	14 to 28	Apply when conditions are favorable for disease development.
Rhizoctonia leaf or sheath spot <i>R. oryzae</i> , <i>R. zea</i>	28	Apply when conditions are favorable for disease development. Rhizoctonia infection can occur under warm, humid conditions on both cool season turfgrass and warm-season turfgrass. This disease has been associated with localized dry spots, and necrotic (brown) ring symptoms can form. The use of a wetting agent can aid in recovery from dry spots.
Rust <i>Puccinia</i> spp.	14 to 28	Apply when conditions are favorable for disease development.
Summer patch <i>Magnaporthe poae</i>	28	Apply when conditions are favorable for disease development. Initiate applications in spring when soil temperatures reach 60° F to 70° F at a 2-inch soil depth, or as dictated by local recommendations. Repeat application at 28-day intervals.
Take-all patch <i>Gaeumannomyces graminis</i> var. <i>avenae</i>	28	Begin applications when conditions are favorable for fungal infection, prior to disease symptom development. Make 2 applications in the fall and subsequent applications in spring depending on local conditions.
Yellow tuft (Downy mildew) <i>Sclerophthora macrospora</i>	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
Zoysia patch <i>Rhizoctonia solani</i> AG2-2LP	14 to 28	Make 1 to 2 applications in the fall when conditions are favorable for disease development, repeat in spring prior to and following greenup. Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.

Application Instructions for Scotts® Pro Granular Fungicide

Scotts® Pro Granular Fungicide is formulated as a dry granule broad-spectrum fungicide which must be broadcast applied with spreader equipment commonly used for ground application. Uniform coverage according to the label rate is essential for disease control. Proper calibration of the spreader equipment is essential before making **Scotts® Pro Granular Fungicide** applications.

Apply with properly calibrated spreader equipment to provide uniform distribution. Check application routinely to determine proper calibration. Avoid overlaps that will increase rates above those specified.

Suggested Spreader Settings For Turf

To apply the label use rate of 3.0 lbs product/1000 sq ft (131 lbs product/acre) and to provide proper product distribution, calibrate spreader according to the settings in the following table. **DO NOT** make more than 5 applications per year at the use rate of 3.0 lbs product/1000 sq ft (131 lbs product/acre).

Spreader	Ground Speed (mph)	Width of Coverage (ft)	Spreader Setting
Professional SS-2 Drop*	3	3	3.5
Turfbuilder® EdgeGuard® DLX (homeowner model)	3	5	3.5
*The Scotts Company does not recommend the use of drop spreaders.			

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of The Scotts Company or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

The Scotts Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE SCOTTS COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND THE SCOTTS COMPANY EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE SCOTTS COMPANY AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

The Scotts Company and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of The Scotts Company.

The Scotts Company
14111 Scottslawn Road
Marysville, OH 43041

©2022 The Scotts Company
All rights reserved