Apheles (Including Sementag) Apheles (Including Sementag) Apheles (Including Sementag) Apheles (Including Sementag) Bits 1, 100 Bits 1, 100 Constructions (Sementage) Bits 1, 100 Bits 1, 100 Constructions (Sementage) Bits 1, 100 Bits 1, 100 Constructions (Sementage) Bits 1, 100 Bits 1, 100 Bit	Sevin® Insect Killer Lawn Granules will provide control of the following pests at the application ra Pest	tes listed below Application Rate
Lond, All, Darly Heb Check, Yaal Hei Charle, Societ Web Check 1 Program	Aphids (including Greenbug) Armyworms (including Lawn Armyworm, Fall Armyworm) Billbugs (including Bluegrass Billbug, Hunting Billbug) Black Turfgrass Ataenius (Adult) Bristletail Caterpillars (including Lucerne Moth) Chinch Bugs Crickets (including House Cricket, Common Short-tailed Cricket, Indies Short-tailed Cricket, Arizona Crickets, Tropical Cricket, Variable Field Cricket, Fall Field Cricket, Southeastern Field Cricket, Texas	
And beelings and beelings an	Cricket, Sand Field Cricket, Vocal Field Cricket, Southern Wood Cricket) Cutworms (including Black Cutworm, Bronzed Cutworm, Granulate Cutworm, Variegated Cutworm) Earwigs Fiery Skipper Firebrat	1 to 2 lbs per 1,000 ft ²
Springene menta were approximately and an approximate of the springene menta were approximately and approximately approximatel	Ladybeetles Leafhoppers Mealybugs Pillbugs Scales (including Ground Pearls) Seed Maggots	
Bernsteinen Meier Der Ausse Jahr Peier Jahr Peier	Springtails Striped Grassworm Two-lined Spittle Bugs Webworms (including Sod Webworm, Grass Webworm, Tropical Sod Webworm) Wireworms	
encode to Dependent and Phasion Ands grouper to Dependent and Phasion Ands grouper to Dependent and Phasion Ands grouper to Dependent and Phasion Science (Dependent Screpton, Leaser Shipptal) Science (Science) The Term of The Term of The Term of Term of Term of Term of Term of Termson, Term of Termson, Termson	Bermudagrass Mites Chiggers Clover Mites Crane Flies (larvae) Ground Beetle Ground Pearl Spider Mites Mole Crickets (including Tawny Mole Cricket, Southern Mole Cricket, Short-Winged Mole Cricket) Spiders (including Brown Recluse Spider, Marbled Cellar Spider, Longbodied Cellar Spider,	2 to 4 lbs per 1,000 ft ²
CPECTED PEST INSTRUCTIONS exploration control or advance and an exploration by applying the post instruction. How and the standards of the instruction of the instruc	Ants (except Carpenter and Pharaoh Ants) European Chafer Fleas (larvae & adults) Grubs mported Fire Ants (adults) (excluding mound treatments) Villipedes Scarab Beetles (includes Japanese Beetle, Green June Beetle, May Beetle) Scorpions (including Arizona Stripetail Scorpion, Southern Unstriped Scorpion, Lesser Stripetail Scorpion, Yellow Ground Scorpion, California Common Scorpion, Northern Scorpion, Eastern	4 lbs per 1,000 ft ²
adybeetie, Twice-stabbed Ladybeetie, Asian Ladybeetie, Maaybug Ladybirdy, Leadhoppers, Lucerne Moth, Meaybugs, Salders (including Jacaness Beetie, Green June Beetie, May Beetie), Scorpions (including Arzona Stripetial Scorpion, Seed Magotis, Sowbugs, Spiders (including Brown Reclues Spider, Marbied Cellar Spic angbodied Cellar Spice, Thintegged Wolf, Thintegged Wolf, Spider, Burnwing Wolf Spider, Prate Wolf Spider, Junne Beetie, Kay Beetie), Scorpions, Tekk et al. Spice, Stripetial Scorpion, Seed Magotis, Sowbugs, Spiders (including Brown Reclues Spider, Marbied Cellar Spice angbodied Cellar Spice, Stripetial Scorpion), Tekk et al. Spices, Spices (Cellar Spice), Spice Marbied Cellar Spice, Spices (Cellar Spice), Spice Marbied Cellar Spice, Spices (Cellar Spice), Spices (Marbied Cellar Spice), Spices (Marbied Spice), Spices (Cellar Spice), Spices (Marbied Cellar Spice), Spices (Marbied Spice), Spice (Marbied Spice), Spices (Marbied Spice), Spice (Marbied Spice), Spice), Spice), Spice (Marbied Spice), Spice), Spice (Marbi	eye appear. This product will provide control of ant mounds within four hours. Food utensils such as teaspoons and nould not be used for food purposes after use with insecticide. For optimum control or for heavily infested areas, cord broadcast treatments. emyourns, Cutworns, and Sod Webworns: Irrigate the treated area with up to 0.1 inches of water immediately at trivtate (release from the granule) the insecticide. Innual Bluegrass Weevil (<i>Hyperodes</i>) adults: Time application for when adult weevils leave their overwintering site rass areas. Ilibug adults: Apply when adult billbugs are first observed during April and May. Iack Turgrass Ataenius adults: Apply during May and July to control the first and second generation of black turing finct Bugs. Trigate the treated area with up to 0.25 inches of water immediately after application to activate (release te insecticide. Higher applications rates may be required to control populations that contain both nymphs and adults armiter. In temperate regions, Spring applications will provide control of over-wintered chinch bugs. Lea larvae: Irrigate the treated area with up to 0.5 inches of water immediately after application to activate (release festicide. Iole Crickets: Apply as late in the day as possible to moist soil (irrigate before application if necessary), and water in cheas of water immediately after treatment. Treat grass areas that receive pressure from adult mole crickets in the si for to peak egg hatch when young nymphs are located near the soil surface. Use higher and more frequent applications to all with yous. Second water applications to activate (release festicide. Iongens: Treat at dusk to prevent outdoor scorpions from entering the building. Treat areas where scorpions resid ay such as in cracks and voids. Remove trash, debris, or firewood that scorpions may use as resting sites prior to the fireks (<i>Including ticks</i> that transmit Lyme Disease and property boundaries where exposure to ticks may occu	I measuring cups Imbine both moun fter application to s and move into rass ataenius adult e from the granule s during the rom the granule) th n with up to 0.5 pring immediately tions to control e hidden during the reatment. juvenile ticks (larva ons (August/Sep- s of home and oth r. Irrigate treated eintroduced from where exposure to at if ticks are ticks tation is known to e performed by important to treat on to ensure they fa fall Armyworm), s. Bristletail, s Short-tailed Texas Field Field Cricket, Jatre Cutworm, Larvae, Fiery d Pearl. Grubs.
FEGETABLES PHI Virichokes 5 Day Steet (Garden) 1 Dag Trassica Vegetables: 7 Day Stroccoli Raab (rapini), Chinese Cabbage (bok choy), Collards, Kale, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens 7 Day Vacuation (Chinese Amsciac Vegetables: 7 Day Stroccoli Rassica Vegetables: 3 Day Strom (sweet) 3 Day Strom (sweet) 3 Day Structurit Vegetables: 3 Day	adybeetle, Twice-stabbed Ladybeetle, Asian Ladybeetle, Mealybug Ladybird), Leafhoppers, Lucerne Moth, J Millipedes, Mole Crickets (including Tawny Mole Cricket, Southern Mole Cricket, Short-Winged Mole Cricket beetles (including Japanese Beetle, Green June Beetle, May Beetle), Scorpions (including Arizona Stripetail S Southern Unstriped Scorpion, Lesser Stripetail Scorpion, Yellow Ground Scorpion, California Common Scor Scorpion, Eastern Sand Scorpion), Seed Maggots, Sowbugs, Spiders (including Brown Recluse Spider, Mar ongbodied Cellar Spider, Shortbodied Cellar Spider, Thinlegged Wolf Spider, Burrowing Wolf Spider, Pirate Spider Mites, Springtails, Striped Grassworm, Ticks (including Brown Dog Tick and ticks that may transmit wo-lined Spittle Bugs, Webworms (including Sod Webworm, Grass Webworm, Tropical Sod Webworm), V Hyperodes and Bluegrass) and Wireworms In the table below, the Pre-Harvest Interval (PHI) refers to the minimum number of days that must pass between st application and the date of harvest. Reapply if insects return, but not more than once every seven days and	Mealybugs,), Pillbugs, Scara Scorpion, pion, Northern bled Cellar Spider Wolf Spider), Lyme Disease), Veevils (including en the date of the
troccoli Raab (rapin), Chinese Cabbage (took choy), Collards, Kale, Mizuna, Mustard Greens, Mustard Spinach, Hape Greens 7 ead & Stem Brassica Vegetables: 7 Broccoli, Chinese Broccoli (gai lon, white flowering broccoli), Brussels Sprouts, Cauliflower, Cavalo Broccolo, Chinese Cabbage (napa) 3 Darm (sweet) 3 3 Structurbit Vegetables: 7 Day Dray otto (Tuil), Chinese Waxgourd (Chinese Preserving Melon), Citron Melon, Cucumber, Gherkin, Gourd (edilble including flyotan, cucuzza, hechima, Chinese orkra), Mormortica spp. (includes balsam apple, balsam par, bitter nelon, Chinese cucumber), Muskmeion, Incore yde melon, Santa Claus melon, San	/EGETABLES Artichokes Beet (Garden) Brassica Vegetables:	PHI5 Days1 Day7 Days
Chayote (fruit), Chinese Waxgourd (Chinese Preserving Melon), Citron Melon, Cucumber, Gherkin, Gourd (edible including hyotan, cucuza, hechima, Chinese orkra), Mormordica spp. (includes balsam apple, balsam pear, bitter nelon, Chinese cucumber), Muskmelon (includes true cantaloupe, casaba, crenshaw melon, golden including hyotan, boney balls, margo melon, Persian melon, pineapple melon, Santa Claus melon, and spaghetti squash), Summer Squash (includes crookneck squash, calabaza, hubbard squash, acorn squash, ind spaghetti squash), Watermelon (includes hybrids and varieties) 3 Day I ead Lettuce 7 Day egume Vegetables - Succulent Edible-Podded Peas, Succulent Shelled Peas and Dried Shelled Peas: 3 Day uward Pea, Edible-pod Pea, Snow Pea, Sugar Snap Pea, Pigeon Pea, English Pea, Garden Pea, Green Pea, Lentil 1 Day upin Sweet Lupin, White Sugar Snap Pea, Norab Bean, Yardlong Bean, Jackbean, Sorab Bean, Wax Bean, Asparagus Bean, Chinese Longbean, Mott Bean, Yardlong Bean, Jackbean, Soybean (immature seed), Swordbean, Lima Bean, Broad Bean (Fava Bean), Blackeyed Pea, Southern Pea, Grain Pea, Grain Pea, Grain Pean, Chinekee Lupin, White Kweet Lupin, Field Bean, Kinder Bean, Aury Kean, Pinto Bean, Chinekee Garbarzo Bean), Guar, Lablab Bean 1 Day ruiting Vegetables (except Sugar Beets): 7 Day variator and Chervil, Chinok, Edible Gana, Gartot, Cassava (Bitter and Sweet), Celeriac (Celery Root), Chavote Root), Radish, Rutabaga, Salsify (Oyster Plant), Black Salsify, Spanish Salsify, Skirret, Sweet Potato, Tamire, Cocoyam), Turmeric, Turnip, Yam Bean, Yam (True) 21 Day Yatus Yama Adama Adama Adama Adamada Ada Pagina Adama Adama Adama Adama Adama Adama Adam	Head & Stem Brassica Vegetables: Broccoli, Chinese Broccoli (gai lon, white flowering broccoli), Brussels Sprouts, Cauliflower, Cavalo Broccolo, Kohlrabi, Cabbage, Chinese Cabbage (napa) Corn (sweet)	7 Days 3 Days
egume Vegetables - Succulent Edible-Podded Peas, Succulent Shelled Peas and Dried Shelled Peas: 3 Day Dwarf Pea, Edible-pod Pea, Snow Pea, Sugar Snap Pea, Pigeon Pea, English Pea, Garden Pea, Green Pea, Lentil 1 Dat Soybean (immature seed), Swordbean, Lima Bean, Broad Bean, Brad Bean, Math Bean, Southern Pea, Grain 1 Dat Soybean (immature seed), Swordbean, Lima Bean, Broad Bean, Brad Bean, Blackeyed Pea, Southern Pea, Grain 1 Dat Garbanzo Bean, Otar, Lablab Bean Free Muthe Lupin, White Sweet Lupin, Field Bean, Kidney Bean, Navy Bean, Pinto Bean, Tepary 21 Dat Garbanzo Bean, Mark Jablab Bean Fruiting Vegetables: 7 Day Roghtan, Okra, Pepino (Melon pear), Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) 7 Day Peanuts 14 Dat Soot and Tuber Vegetables (except Sugar Beets): 14 Dat Arraku, Tarier, Cocoyam), Turmeric, Turnip, Yam Bean, Yam (True) 21 Dat Varsley, Parsnip, Oriental Radish (Daikon), Radish, Rutabaga, Salisty (Oyster Plant), Black Salisty, Spanish Salisty, Skirret, 21 Dat Soybean 3 Dat Sophean 3 Dat	Chayote (fruit), Chinese Waxgourd (Chinese Preserving Melon), Citron Melon, Cucumber, Gherkin, Gourd (edi including hyotan, cucuzza, hechima, Chinese orkra), <i>Mormordica</i> spp. (includes balsam apple, balsam pear, l nelon, Chinese cucumber), Muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, gold pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus n ind snake melon), Pumpkin, Summer Squash (includes crookneck squash, scallop squash, straightneck squa regetable marrow, zucchini), Winter Squash (includes butternut squash, calabaza, hubbard squash, acorn squ ind spaghetti squash), Watermelon (includes hybrids and varieties)	bitter en 3 Days nelon, ish, iash,
Rumer Bean, Snap Bean, Wax Bean, Asparagus Bean, Chinese Longbean, Moth Bean, Yardlong Bean, Jackbean, Soybean (immature seed), Swordbean, Lima Bean, Broad Bean (Fava Bean), Blackeyed Pea, Southern Pea, Grain Lupin, Sweet Lupin, White Sweet Lupin, Field Bean, Xidney Bean, Navy Bean, Pirot Bean, Tepary Bean, Adzuki Bean, Catjang, Cowpea, Crowder Pea, Moth Bean, Mung Bean, Nice Bean, Urd Bean, Chickpea1 Dag (Succult 21 Day 20 Day 21 Day 21 Day 22 Day 23 Day1 Dag 23 Day 24 Day 24 Day 24 Day 24 Day 24 Day 24 Day1 Day 24 Day 25 Day 26 D	Legume Vegetables - Succulent Edible-Podded Peas, Succulent Shelled Peas and Dried Shelled Peas: Dwarf Pea, Edible-pod Pea, Snow Pea, Sugar Snap Pea, Pigeon Pea, English Pea, Garden Pea, Green Pea, Len	(Dried)
Figplant, Okra, Pepino (Melon pear), Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper)7 DayPeanuts14 DayRoot and Tuber Vegetables (except Sugar Beets): Arracacha, Arrowroot, Edible Burdock, Edible Canna, Carrot, Cassava (Bitter and Sweet), Celeriac (Celery Root), Chayote (Root), Turnip-Rooted Chervil, Chicory, Chufa, Dasheen (Taro), Ginger, Ginseng, Horseradish, Leren, Turnip-Rooted Parsley, Parsnip, Oriental Radish (Daikon), Radish, Rutabaga, Salsify (Dyster Plant), Black Salsify, Spanish Salsify, Skirret, Weet Potato, Tanier (Cocoyam), Turmeric, Turnip, Yam Bean, Yam (True)21 DayPotato21 DaySoybean21 DaySoybean21 DaySignach40 DayImage: RENES, FRUITS, AND NUTSPHIBlueberries1 DaySapheerries including: Blackberries, Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries, Blackberries Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries, Brapes30 Day Tree Nuts Group including: Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia ro but, Pistachio, Walnut (Black and English)7 Day	Runner Bean, Snap Bean, Wax Bean, Asparagus Bean, Chinese Longbean, Moth Bean, Yardlong Bean, Jackbe Soybean (immature seed), Swordbean, Lima Bean, Broad Bean (Fava Bean), Blackeyed Pea, Southern Pea, Gr Lupin, Sweet Lupin, White Lupin, White Sweet Lupin, Field Bean, Kidney Bean, Navy Bean, Pinto Bean, Tepary Bean, Adzuki Bean, Catjang, Cowpea, Crowder Pea, Moth Bean, Mung Bean, Rice Bean, Urd Bean, Chickpea (Garbanzo Bean), Guar, Lablab Bean	an, 1 Day ain (Succulen 21 Days (Dried)
Sweet Potato, Tanier (Cocoyam), Turmeric, Turnip, Yam Béan, Yam (True) 21 Day Potato 21 Day Soybean 21 Day Spinach 40 Day formato 1 Day Blueberries 1 Day Caneberries including: 1 Day Blackberries, Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries, 3 Day 30 Day Grapes 30 Day Tree Nuts Group including: 30 Day Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia rub, Pistachio, Walnut (Black and English) 7 Day	Eggplant, Okra, Pepino (Melon pear), Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepp Peanuts Root and Tuber Vegetables (except Sugar Beets): Arracacha, Arrowroot, Edible Burdock, Edible Canna, Carrot, Cassava (Bitter and Sweet), Celeriac (Celery Root), Cha Root), Turnip-Rooted Chervil, Chicory, Chufa, Dasheen (Taro), Ginger, Ginseng, Horseradish, Leren, Turnip-Rooted	14 Days
Billeberries PHI Blackberries including: 1 Day Blackberries, Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries, 3 Day Blackberries 30 Day Grapes 30 Day Tree Nuts Group including: 30 Day Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia 7 Day nut, Pistachio, Walnut (Black and English) 7 Day	Sweet Potato, Tanier (Cocoyam), Turmeric, Turnip, Yam Bean, Yam (True) Potato Soybean Spinach	Skirret, 21 Days 21 Days 40 Days
Caneberries including: 3 Day Blackberries, Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries, 3 Day Raspberries 30 Day Grapes 30 Day Tree Nuts Group including: 3 Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia 7 Day nut, Pistachio, Walnut (Black and English) 7		4.5
Grapes 30 Day Tree Nuts Group including: Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia 7 Day nut, Pistachio, Walnut (Black and English) 7 7		
nut, Pistachio, Walnut (Black and English)	<mark>Blueberries</mark> Caneberries including: Blackberries, Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries	PHI 1 Day
	Blueberries Caneberries including: Blackberries, Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries Raspberries Grapes Free Nuts Group including:	PHI 1 Day 3 Days 30 Days










Net Weight: 20 LB. (9.07 kg)





[‡]When used as directed








GardenTech is a registered trademark of Gulfstream Home & Garden, Inc Sevin is a registered trademark of Tessenderlo Kerley, Inc.

Scotts, Turf Builder and Edgeguard are registered trademarks of OMS Investments, Inc. Sta-Green is a registered trademark of LF, LLC.



TRUSTED BRAND FOR OVER 50 YEARS

QUICK FACTS

Ils lawn-damaging insects that cause your lawn to thin and turn brown

ects edibles from insect damage

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent permitted by applicable law, User agrees to hold Seller harmless for any claims relating to such factors. LAWN, PERIMETER, and GARDEN APPLICATIONS

Broadcast Sevin® Insect Killer Lawn Granules to provide uniform coverage over the treatment area. Locate the listed pest to be controlled in the following table and follow the specified application rate to provide excellent control under typical conditions. When treating for multiple pests, apply the higher of the specified rates. Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed insect activity. Repeat application no more than once per 7 days. Do not exceed 4 lb per 1,000 ft² per application. Best results are obtained if the treated area is thoroughly watered immediately after application. Residential Lawn Application: Broadcast this product with spreader equipment to ensure uniform coverage over the treatment area. Perimeter Application: Apply to ground area 5 to 10 feet wide around and adjacent to the structure. **Ornamental & Flower Garden Application:** Use this product as a spot treatment or area treatment for pest infestations on or around ornamental trees, shrubs, flowers and ground covers. Certain plants such as annuals may be sensitive to the watered-in granules. A limited area should be treated and observed for one week prior to application to the entire planting. Fruit & Vegetable Garden Application: Apply this product to control listed pests in home fruit and vegetable gardens either as a preventative measure or to control existing infestations. Preventative Applications: This product may be applied just prior to planting or after plants emerge and worked into the top 4 to 6 inches of soil. Use 1-4 lb of product for a 1,000 sq ft area and immediately water the area thoroughly after application for best results.

Applications to Control Existing Infestations: Apply this product to the garden area using 1 to 4 lb of product for a 1,000 sq ft area. Best results are obtained if the treated area is thoroughly watered immediately after application. Treat when insects first appear and retreat as necessary. SPREADER EQUIPMENT SETTINGS Broadcast application should be performed with a calibrated broadcast spreader to evenly distribute the granules. Approxi-mate settings for several models are provided in the table below. Mow lawn first for best results. SPREADER SETTINGS - -

SPREADER SETTINGS			NOTE: Spreader settings are approximate. It is
Broadcast Spreaders	Rate (lbs/1,000 ft ²)	Setting	recommended that you calibrate your spreader before
Scotts [®] Turf Builder [®] Edgeguard [®] DLX	1 2 4	4 1/2 7 12	application since manufacturer's settings can vary among the same model of spreader. Broadcast Spreader: If your lawn is rectangular, apply two
Scotts [®] Turf Builder [®] Edgeguard [®] Mini	1 2 4	3 3/4 5 1/2 12	border strips across each end for a turning area. If your lawn is an irregular shape, apply a border strip completely around it. Then apply product back and forth in the longest
Sta-Green [®] Rotary	1 2 4	4 3/4 9	direction. Turn the spreader off and on at the edge of each strip to prevent uneven application. Overlap the throw pattern by about a foot to ensure complete coverage.
Drop Spreaders	Rate (lbs/1,000 ft ²)	Setting	Drop Spreader: Apply granules with spreader going in the same direction as the longest part of the lawn. Apply two (2)
Scotts [®] Turf Builder [®] Edgeguard [®] Classic	1 2 4	5 1/4 8 10	header strips first. To ensure thorough coverage, make sure to overlap wheel marks. Shutoff the spreader when finished or when the header strips have been reached.
	EPA Est. 6	7572-GA	3346-71004 \-1 _A , 67572-GA-2 _B irst letter of run code on container.

6 **13499 0102**

RT051617AAAY

Distributed by: TechPac, L.L.C. 1000 Parkwood Circle, Suite 700 Atlanta, GA, 30339