

FRONT



→
WEB DIRECTION

5.59"(h) x 3.38"(w)

BOOKLET OUTSIDE



The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

	Description: Daconil 16oz Back ECL		NOTES Measurements are in inches.	LEGEND Copy Image Area Base Cut/Fold Lines Book Cut/Fold Lines
	Supplied for: Central Garden & Pet	File No.: T00634		
	Finished Size: 5.59" (H) x 3.062" (W)	Supersedes:		
	Date: 6 / 26 / 15	Supplied By:		
Application: Illustrator CS6	Prepared By: BM			

LEFT

2.312 in

5.590

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Shake Well Before Using

Read Entire Label and Use Strictly in Accordance with Label Precautionary Statements and Directions.

For any questions regarding disease identification, consult an extension agent, university or local arborist.

tsp = teaspoon Tbs = Tablespoon

Readily mixes with water and can be used in a hand-held, hose-end or tank-type sprayer. Shake well before using.

Fine spray required for proper coverage; use a hose-end sprayer for best results.


Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high schools), campgrounds, churches and theme parks. Do not apply this product in a way that will contact other persons or pets, either directly or through drift. Keep people and pets off treated areas until dry.

Vegetable Diseases

Apply the amount of concentrate in the table in one gallon of water to cover 200 sq ft (10' x 20').

Thoroughly spray to cover all plant surfaces. **Note:** Avoid spraying plants during extremely hot and sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to animals.

RIGHT



Crop	Diseases Controlled	Amount of Concentrate	Application Notes	PHI*
Beans (Green)	Rust, Botrytis Blight (Gray Mold)	1 Tbs	Start spraying at first sign of disease. Apply every 7 days to maintain control.	7
Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Alternaria Leaf Spot, Downy Mildew	1 1/2 tsp	Start spraying when conditions favor disease, after transplanting or when sprouts appear. Apply every 7 to 10 days.	0
Brussels Sprouts (only in CA)	Ring Spot	2 tsp	Start spraying at first sign of disease. Apply every 7 to 10 days to maintain control.	0
Carrots	Cercospora (Early) Blight, Alternaria (Late) Blight	2 tsp	Start spraying after transplanting. Apply every 7 days.	7
Celery	Cercospora (Early) Blight, Septoria (Late) Blight, Basal Stalk Rot, Pink Rot (suppression)	1 1/2 tsp	Start spraying when conditions favor disease. Apply every 7 to 10 days to maintain control. Do not apply to corn if you plan to process it.	14
Corn (Sweet)	Helminthosporium Leaf Blight, Rust	1 Tbs	Start spraying when plants are in the first true leaf stage or when conditions favor disease. Apply every 7 days to maintain control.	0
Cucumbers, Melons, Pumpkins, Squash	Anthraxnose, Downy Mildew, Target Spot, Scab, Gummy Stem Blight (Black Rot), Leaf Blight, Cercospora Leaf Spot, Powdery Mildew (except Southwestern states)	2 tsp	Apply every 7 days to maintain control.	0

(continued)

*PHI = the days required between last application and harvest.

LEFT

5.590

2.312 in

Crop	Diseases Controlled	Amount of Concentrate	Application Notes	PHI*
Cucumbers	Fruit Rot (suppression)	1 Tbs	Start spraying when plants are in the first true leaf stage or when conditions favor disease. Apply every 7 days to maintain control.	0
Onion (Green), Leek, Shallot	Botrytis Leaf Blight (Blast), Purple Blotch, Downy Mildew (suppression)	1 Tbs	Start spraying at first sign of disease. Apply every 7 to 10 days. Do not use more than 3 times per year.	14
Onion (Dry Bulb), Garlic			Start spraying at first sign of disease. Apply every 7 to 10 days.	7
Potato	Early Blight, Late Blight, Botrytis Vine Rot	1 1/2 tsp	Start spraying when plants are 6 inches high or at first sign of disease. Apply every 5 days to maintain control.	0
Tomato	Early Blight, Late Blight, Gray Leaf Spot, Gray Leaf Mold, Septoria Leaf Spot, Anthracnose, Alternaria Fruit Rot (Black Mold), Rhizoctonia Fruit Rot, Botrytis Gray Mold	1 Tbs	Start spraying when conditions favor disease. Apply every 7 to 10 days to maintain control.	0

(continued)

*PHI = the days required between last application and harvest.

RIGHT

5.590

2.312 in

Fruit Diseases
Apply the amount of concentrate in the table below in one gallon of water. Thoroughly spray to cover all plant surfaces.

Fruit Tree	Diseases Controlled	Amount of Concentrate	Application Notes
Peach, Nectarine, Apricot	Peach Leaf Curl, Coryneum Blight (Shot-hole)	2 1/4 tsp	Do not apply between fruit formation and harvest. Spray at leaf fall in late autumn. Reapply 1 to 2 more times in mid- to late winter before buds begin to swell. For Shot-hole, to prevent fruit infections, also apply 1 to 2 weeks after petal fall when fruit just begins to form.
	Brown Rot, Blossom Blight, Scab		
Cherry, Plum, Prune	Brown Rot Blossom Blight, Cherry Leaf Spot (cherries)	2 1/4 tsp	Do not apply between fruit formation and harvest. Spray when buds begin to show color. Reapply at full bloom. If cool, moist weather continues, make a third application at petal fall. For Cherry Leaf Spot, also apply when fruit just begins to form and again 7 days after fruit is removed; apply once more 10 to 14 days later, if tree has a history of leaf spot.
	Alternaria Fruit Spot		
Papaya	Alternaria Fruit Spot	1 1/2 tsp	Start spraying when fruit spots appear (April to July). Spray every 14 days until dry weather returns.
Passion Fruit (only in HI)	Anthracnose, Stem End Rot	3/4 tsp	Start spraying before spots appear on fruit (April to July). Spray every 14 days until dry weather returns.
	Alternaria Fruit and Leaf Spot (Passion Fruit Brown Spot)		

LEFT

5.590

2.312 in

Unless otherwise directed in the table below, mix 2 1/4 tsp per gallon of water or 3 Tbs (1/2 oz) per 4 gallons of water and spray foliage to point of runoff. Begin applications as directed for each species and disease condition noted and apply every 7 to 14 days (unless otherwise specified) until conditions no longer favor disease. During periods when conditions favor severe disease, generally cloudy or wet weather, use the shorter interval between applications.

Species	Diseases Controlled	Suggested First Application
Broadleaf Shrubs and Trees		
Ash	Cercospora, Cercosporium, Cylindrosporium, Leaf Spots	Spring bud break
Azalea*, Rhododendron*	Phytophthora Die-Back, Ovulinia Flower Blight	New leaf emergence; early bloom
Buckeye (Horseshoe)	Leaf Blotch, Anthracnose	Spring bud break
Crabapple	Scab, Cedar-apple Rust, Sphaeroopsis, Leaf Spot	Spring bud break
Dogwood	Anthracnose (Discula sp.), Septoria Leaf Spot	Spring bud swell; early bloom
Eucalyptus	Anthracnose	Spring bud break
Firethorn (Pyracantha)	Scab	Spring bud break
Flowering Almond, Quince and Cherry Hawthorn	Monilia Blossom/Branch Blight, Rust, Fabraea Leaf Spot	Early bloom; pre-bloom
Holly	Rhizoctonia Web Blight	Warm, moist conditions
Mountain Laurel	Cercospora Leaf Spot	Spring bud break
Oak (only red group)	Taphrina Blister, Actinopelle Leaf Spot, Anthracnose	Dormant bud swell
Oregon-Grape (Mahonia)	Rust	Spring bud break

(continued)

RIGHT

5.590

2.312 in

Species	Diseases Controlled	Suggested First Application
Broadleaf Shrubs and Trees - continued		
Photinia	Fabraea (<i>Entomosporium</i>) Leaf Spot	Spring bud break
Pieris (Andromeda)	Phytophthora Die-Back	New leaf emergence
Poplar	Marcosonia Leaf Spot	Spring bud break
Privet	Cercospora Leaf Spot	Prolonged wet conditions
Sycamore (Plane tree)	Anthracoze	Spring bud break
Viburnum	Powdery Mildew	Mid-summer
Bulbs and Flowering Plants		
Carnation	Alternaria Leaf Spot, Branch Rot, Botrytis	Transplant of cuttings; cool, moist conditions
Chrysanthemum (Daisy)	Mycosphaerella Ray Blight, Septoria Leaf Spot	Transplant of cuttings
Geranium	Botrytis Flower Blight (Gray Mold), Rust	Pre-bloom; cool, moist conditions
Gladiolus	Curculiaria Leaf Flower Spot, Botrytis Leaf/Flower Spot	Early propagation
Hollyhock	Rust	Early seedling stage
Hydrangea* (only foliage)	Cercospora and Septoria Leaf Spots, Rust	Early propagation
Iris	Botrytis Blossom Blight, Diymellina Leaf Spot	Cool, moist conditions
Lily	Botrytis Gray Mold (folar phase), Phytophthora Blight	Pre-bloom

(continued)

*Discoloration of blooms has been noted on certain varieties when applications are made during flowering.

LEFT


5.590

2.312 in

Species	Diseases Controlled	Suggested First Application
Bulbs and Flowering Plants - continued		
Petunia*	Botrytis Gray Mold (inlar phase), Phytophthora Blight, Botrytis Blight	Pre-bloom
Rose—Use 1 1/4 fl oz (2.12 Tbs) per 4 gallons of water	Black Spot, Botrytis Blight	Spring bud break
Statice	Anthraxose, Cercospora, Alternaria, Leaf Spots, Botrytis, Leaf Blight	Spring bud break
Zinnia	Powdery Mildew	First sign of disease
Syringium	Cephalosporium Leaf Spot	Warm, moist conditions
Foliage Plants		
Dracaena	Fusarium Leaf Spot	Pre-transplant
Prachysandra — Use 5 tsp per gallon of water	Volutella Leaf Blight	Spring bud break
Leatherleaf fern	Ascochyta Blight, Cercospora Leaf Spot, Cythrodactylum Leaf Spot, Rhizoctonia Blight	Spring bud break
Prayer Palm (Chamaedorea)	Bipolaris Leaf Spot (<i>Helminthosporium</i>)	Cool, moist conditions
Prayer Plant (Maranta)	Helminthosporium Leaf Spot	Early propagation
Oyster Plant (Rhoeo)	Tan Leaf Spot	Early propagation
Philodendron	Phytophthora Blight, Dactyaria Leaf Spot	Moist conditions
Syringium	Cephalosporium Leaf Spot	Warm, moist conditions

*Discoloration of blooms has been noted on certain varieties when applications are made during flowering.

RIGHT



LEFT

2.312 in

5.590

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry (preferably locked) place inaccessible to children and pets.

Pesticide Disposal: Nonrefillable container; do not reuse or refill this container. **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Disposal: Triple rinse (or equivalent) container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap; shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Buyer assumes all risk of use, storage or handling of this product not in strict accordance with this label.

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347.

If you have questions relating to product use or application, please call (866) 945-5033.

If you have a product spill or transportation emergency call (800) 424-9300.

GardenTech is a registered trademark of Gulfstream Home & Garden, Inc. Daconil is a registered trademark of GB Biosciences Corp.

BASE LABEL

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

WARNING: Causes substantial but temporary eye irritation. Causes skin irritation. Harmful if inhaled, swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse. Wear goggles or face shield, coveralls over a short-sleeve shirt and pants, and rubber gloves when mixing and applying this product.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash in exposed skin areas. Persons having an allergic reaction should contact a physician.

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 800-420-9347 for emergency medical treatment information.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If Swallowed: Immediately call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Environmental Hazards

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. This chemical can contaminate surface water through spray drift. Under some conditions it may also have high potential for runoff into surface water for several days to weeks after application; these include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow groundwater, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips and areas overlaying tile drainage systems that drain to surface water.

EPA Reg. No. 67572-82-71004
EPA Est. 67572-GA-1_a, 67572-GA-2_b
Subscript used is first letter of run
code on container.

Distributed by:
TechPac, LLC,
1000 Parkwood Circle
Suite 700
Atlanta, GA 30339

file: 68658_base_art
folder: 68658_Central G&P_Daconil
color: black, 186
dp proofed _____

created by: customer
alt: 10-30-15 jw
size: 3.062"(w) x 5.59"(h)