

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions, use restrictions and use precautions.

HISE RESTRICTIONS.

- •This product may only be applied against the pests listed on this label, at the use sites listed on this label, and by the application methods listed on this label.
- memoralised on mis label.

 This profluct may only be used to control pocket gophers** and moles* in manual, below-ground applications in lawns.

 Bait must be applied directly into the burrow or tunnel systems.
- Do not dpply in gardens and areas where food or feed may become contaminated.

 Any person who retrieves carcasses or unused bait following
- application of this product must wear waterproof gloves.

 Do not apply this product by the use of an artificial burrow builder.
- Not to be sold in individual containers containing more than 1 lb. of

APPLICATION DIRECTIONS FOR POCKET GOPHERS**: Apply bait to control packet gophers** using one of the two methods outlined

Plug Rembval Method/Baiting:
Burrowing pocket gophers** (Thomomys spp. and Georthys spp.)
throw out low, fars-staped mounds on either side of their underground trunnel. These lateral humes Coming to the surface are on the flat side
of the find and are plugged with soil. Using a long-handled spoon, carefully remove the plug. Insert one teaspoon of bait as fal down as possible. Reclose opening, being careful not to cover bait with soil.



88603 03722

1005

- Not for mole use in California or Indiana. Not for use against star-nosed mole in North Carolina.

 Not for use against gophers in North Carolina.

Questions or Comments? all 1-877-332-0755 www.tomcat.com Available Exclusively From: Tomcat Brands Marysville, OH 43040 © 2016 Tomcat Brands. World rights res Main Runway Method/Baiting: To locate main runways, probe about 6 to 8 inches away from mounds by use of an iron bar, a strong stick, or other suitable implement. Resistance on probe will drop suddenly and probe will fall about two inches when main runway is entered. Carefully enlarge the opening to accommodate insertion of the bait. Insert one teaspoon of bait into each active tunnel. Carefully close the hole with a stone, sod, or soil. Take care not to cover bait with soil.

APPLICATION DIRECTIONS FOR MOLES*: The presence of moles may be indicated by a network of surface ridges in the turf, and/or by a series of conical mounds of earth pushed up from deep burrows.

Surface Ridges:

Prior to treatment, determine which burrows are active according to one of the following two methods. Either press down a small section of the burrow or remove a one-inch section of the burrow's roof. Mark these sections and recheck in 24 hours. The burrow is considered active if the flattened burrow has been raised up or if the roof has been repaired. Only treat active burrows. Carefully punch a hole in the top of an active tunnel, drop a teaspoonful amount of this product into the burrow and carefully close the hole with stone or sod. Do not collapse the tunnel again or allow loose soil to cover the bait. After a few days check to see if the moles are still active as described under "Surface Ridges" above. If so, repeat treatment.

Conical Mounds:

Remove center plug from the conical mounds with a long-handled spoon and drop a teaspoonful amount of this product into the underground runway. Close opening with stone or sod. Do not allow

FOLLOW-UP: Wearing waterproof gloves, collect and dispose of dead, exposed onimals. Dispose of leftover bair in accordance with disposal instructions. Use detergent and hot water to wash spoons used for application into hurrows. Do not use spoon for mixing holding or transferring food or feed. If reinfestation does occur repeat treatment.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store only in original container in a dry

PESTICIDE DISPOSAL AND CONTAINER HANDLING:
Non-refillable container. Do not reuse or refill this container. If empty: Place in trosh or offer for recycling if outsideble. If partially filed: Place in trosh or cell your local solid waste agency for disposal instructions. Never place unused product down any independent of the partially filed: Place in trosh or cell your local solid waste agency. indoor or outdoor drain.

Notice: To the extent consistent with applicable law, buver assumes all risks of use, storage or handling of this product not in accordance with directions.

Tomcat Brands Guarantee: If for any reason you, the consumer, are not satisfied with this product, mail us your original proof of purchase to obtain a full refund of your purchase price.

102715/07-15

CAUTION: Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soop and water offer handling and before earing, diming, chewing gum, using tobocco, or using the tolet. Wear long sleeked shirt and long barnts, shoes plus socks and gloves. Remove and wesh contaminated clothing before reuse. Any person with retrieves corcesses or unused bait following application of this product must

FIRST AID

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-332-0755 for emergency medical treatment information.

If you experience signs and symptoms such as nausea, abdominal pain, tightness in chest, or weakness, see a physician immediately.

 Call a poison control center or immediately for treatment advice or t f swallowed the patient to the nearest hospital.

If in eves

If inhaled

 Do not drink water. Do not administer anything by mouth or induce vomiting unless told to do so by the poison

transport

- control center or doctor. • Hold eye open and rinse slowly and gently with
- water for 15–20 minutes.

 Remove contact lenses, if present, after the first
- five minutes, then continue rinsing eye.

 Call a poison control center or doctor for
- treatment advice. · Move person to fresh gir.

If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice.


TREATMENT FOR PET POISONING

NOTE TO PHYSICIAN OR VETERINARIAN

Contains the phosphine-producing active, Zinc Phosphide.
Probable mucosal damage may contraindicate the use of gastric Produce incusts admigs may communicate the use or gistin. Growbe. For animals ingesting bid and/or showing poisoning signs, induce vamiting by using hydrogen peroxide. Sodium bicalbonate can be given orally to neutralize the stomach acidity. The Istomach and intestinal tract can be evacuated, oxygen administered and cardiac and circulatory stimulants given.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to birds, fish and other wildlife. Wildlife feeding on treated bait may be killed. Dogs, cats, and other



KEEP OUT OF REACH OF CHILDREN CAUTION

See back and side panels for First Aid, ACTIVE INGREDIENT: Zinc Phosphide (CAS#1314-84-7) OTHER INGREDIENTS: TOTAL additional precautionary statements, and Directions for Use. EPA REG. NO. 90780-4 EPA EST. NO. 12455-WI-1

NET WT. **6 OZ** (168 g)