
REFILL
KILLS
CRABGRASS
PPN 10011U1-06 LB15627

Roundup®
FOR LAWNS,

ACTIVE INGREDIENTS:

MCPA, dimethylamine salt	0.257%
Quinclorac	0.118%
Dicamba, dimethylamine salt	0.029%
Sulfentrazone	0.015%
OTHER INGREDIENTS:	99.581%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

0.018 lb 2-methyl-4-chlorophenoxyacetic acid equivalent per gallon or 0.210%
0.010 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 0.118%
0.002 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.024%
0.001 lb N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.015%

KILLS WEEDS · WON'T HARM LAWNS*

UP TO 3 MONTH CONTROL**

KILLS CRABGRASS · DANDELION · CLOVER · YELLOW NUTSEDGE***

FAST ACTING FOR VISIBLE RESULTS IN HOURS

Keep Out Of Reach Of Children

CAUTION Stop! Read the entire label first. Observe all precautions and follow directions carefully. See back panel booklet for additional precautionary statements and first aid.

Mantener Fuera Del Alcance De Los Niños

PRECAUCIÓN ¡Alto! Lea primero toda la etiqueta. Observe todos los avisos de precaución y siga atentamente las instrucciones. También preste atención a los avisos de precaución y los primeros auxilios en la contraportada.

PPN 10011U1-04

LB15468

NET CONTENTS / CONTENIDO NETO: 1.25 GALLONS (1 GAL 32 FL OZ)

* When used as directed
** As listed
*** See inside booklet for weed list

OPEN
/ ABRA

READY-TO-USE

FOR USE ON NORTHERN GRASSES

KILLS 253 TYPES OF WEEDS**

ONE STEP WEED CONTROL FOR UP TO 3 MONTHS



For best results in non-lawn areas like garden beds, patios and driveways, use Roundup® Ready-To-Use Weed & Grass Killer III

† **GUARANTEED RESULTS:** If for any reason you are not satisfied after using this product, simply send us original proof of purchase and we will refund the purchase price.

Questions & Comments Preguntas y comentarios
Call 1-800-246-7219 or Visit our Website at www.roundup.com

Produced for:
Monsanto Company
Lawn & Garden Products
800 N. Lindbergh Blvd.
St. Louis, MO 63167
© 2020 Monsanto Company

EPA Reg. No. 2217-917-71995
EPA Est. No. 239-IA-3

** As listed

PPN 10011U1-05
LB15469

1005

FPO

0 70183 50201 1

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RESTRICTIONS:



- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Do not allow people or pets to enter treated area until spray has dried.

PRECAUTIONS:

Roundup® For Lawns$_1$ is a ready-to-use (RTU) product intended for "spot" treatment of individual broadleaf and grass weeds by homeowners on residential lawns. This product may be applied to lawns containing the following species of grass: Kentucky bluegrass, perennial ryegrass, fescue species (including tall, red, and fine leaf varieties), bermudagrass[1], buffalograss and zoysiagrass.

[1] Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

Roundup® For Lawns$_1$ should not be used on lawns containing bentgrass, St Augustinegrass, bahiagrass, centipedegrass, or carpetgrass. Do not use on dichondra or lawns with desirable clovers or legumes, on vegetable gardens, fruits, or ornamentals (flowers, trees, groundcovers, hedges, landscape/flower beds and shrubs).

WEEDS CONTROLLED:

Roundup® For Lawns$_1$ will control or suppress the following broadleaf and grass weeds.

BROADLEAF WEEDS

Amaranth, purple
Amaranth, slender
Aster, heath
Aster, white heath
Aster, purple
Aster, white prairie
Artichoke
Austrian fieldcress
Bedstraw, catchweed
Bedstraw, smooth
Bedstraw, northern
Bedstraw, pineywoods
Beggarticks, hairy
Betony, Florida
Bindweed, field
Bindweed, hedge
Bitter Wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black medic
Blood flower milkweed
Blue lettuce
Brassbuttons
Bristly oxtongue
Broomweed
Buckhorn
Bullnettle
Burclover
Burdock
Burweed, lawn (spurweed)
Buttercup, bulbous
Buttercup, creeping
Buttonweed, common (poorjoe)
Buttonweed, Virginia
Carolina geranium (wild)
Carpetweed
Catnip
Catsear, spotted (false dandelion)
Chamber bitter
Chickweed, common
Chickweed, mouseear
Chickweed, sticky
Chicory
Cinquefoil
Clover, crimson
Clover, alsike
Clover, hop
Clover, rabbit foot
Clover, red
Clover, strawberry
Clover, yellow sweet
Clover, white sweet
Clover, white
Cockle
Cocklebur
Cocklebur, oriental
Cocklebur, spiny
Common mullein
Corn spurry
Creeping beggarweed
Creeping Jenny (creeping Charlie)
Cupid shaving brush
Cudweed
Daisy, English
Daisy, oxeye
Dandelion, common
Dandelion, false
Day flower
Deadnettle
Dichondra
Dock, broadleaf
Dock, curly
Dogbane
Dogfennel
Dollarweed (pennywort)
Doveweed
Elderberry
Falseflax
False sunflower
Fiddleneck
Filaree, redstem
Filaree, whitestem
Fleabane, daisy (annual)
Fleabane, rough
Galinsoga, hairy
Galinsoga, smallflower
Goldenrod
Ground ivy
Groundsel
Gumweed
Hairy fleabane
Hawkweed, meadow
Hawkweed, mouseear
Hawkweed, orange
Hawkweed, yellow
Healall
Heartleaf drymary
Hemp
Henbit
Hoary cress
Horsenettle
Horseweed
Jimsonweed
Johnny-Jumpup violet
Knawel
Knotweed, prostrate
Kochia
Lambsquarters
Lespedeza
Mallow, alkali
Mallow, bristly
Mallow, common
Mallow, dwarf
Mallow, Venice
Marcela

Matchweed
Mexicanweed
Moneywort
Morningglory, bigroot
Morningglory, ivyleaf
Morningglory, Mexican
Morningglory, red
Morningglory, tall
Morningglory, woolly
Mugwort
Mustard, ball
Mustard, black
Mustard, blue
Mustard, hedge
Mustard, Indian,
Mustard, tansy
Mustard, tumble
Mustard, white
Mustard, wormseed
Nettle, stinging (burning)
Nettle, tall
Nettle, wood
Parsley-piert
Parsnip
Pearlwort
Pennycress (Frenchweed)
Pepperweed, perennial
Pepperweed, Virginia (peppergrass)
Pigweed, prostate
Pigweed, redroot
Pigweed, Russian
Pigweed, smooth
Pigweed, tumble
Pineappleweed
Plains coreopsis (tickseed)
Plantain, blackseed
Plantain, bracted
Plantain, broadleaf
Plantain, buckhorn
Plantain, hoary
Plantain, narrowleaf
Plantain, slender
Plantain, woolly
Poison ivy
Poison oak
Pokeweed
Prairie sunflower
Prickly lettuce (compass plant)
Prickly sida
Prostrate knotweed
Puncturevine
Purslane, common
Pusley, Brazil
Pusley, Florida
Pusley, large flower
Ragweed, bur
Ragweed, common
Ragweed, lanceleaf
Ragweed, western
Redstem filaree
Rough cinquefoil
Scarlet pimpernel
Shepherdspurse
Smartweed, ladysthumb
Smartweed, pale
Smartweed, Pennsylvania
Smooth chaff-flower
Smooth dock
Sorrel, red (sheep)
Sowthistle, annual
Sowthistle, spiny
Spanish needles
Speedwell, bi-lobed
Speedwell, common
Speedwell, corn
Speedwell, ivyleaf
Speedwell, Persian
Speedwell, purslane
Speedwell, slender
Speedwell, snow
Speedwell, thymeleaf
Spiny amaranth
Spurge, spotted
Spurge, prostrate
Spurweed
Star of Bethlehem
Strawberry, Indian mock
Tansy ragwort
Tanweed
Thistle, blessed
Thistle, bull
Thistle, Canada
Thistle, flodman (prairie)
Thistle, musk
Thistle, platte
Thistle, plumeless
Thistle, Russian
Thistle, Scotch
Thistle, wavyleaf
Thistle, yellow (horrible)
Trailing crownvetch
Velvetleaf
Vervain, blue
Vervain, hoary
Vervain, prostrate
Vervain, tall
Vetch, bird
Vetch, hairy
Vetch, milk
Vetch, narrowleaf
Vetch, wild
Virginia creeper
Western clematis
Western salsify
Wild aster
Wild buckweat
Wild carrot
Wild four-o-clock
Wild garlic
Wild lettuce
Wild marigold
Wild mustard (yellow rocket)
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild violet
Woodsorrel, creeping
Woodsorrel, violet
Woodsorrel, yellow (Oxalis)
Woolly croton
Wormseed
Yarrow, common
Yarrow, western
Yarrow, yellow

GRASSY WEEDS

Crabgrass (large and smooth)
Foxtail (green, giant and yellow)
Yellow nutsedge

WHEN TO USE

Apply this product to actively-growing weeds during the early growth stages for the best results. Two or more applications at 14 day intervals throughout the season may be required for grass weeds. Apply when daytime temperatures are between 45° and 90° F.

SPRING AND EARLY SUMMER:

- **Crabgrass (large and smooth):** Apply this product from crabgrass emergence until the 6 leaf stage (2 tillers) or when the crabgrass is 3 to 4 inches tall. This period begins the first time that you can see crabgrass in your lawn. Use two applications with a 14-day interval between applications.
- **Foxtail (green, giant and yellow):** Apply this product to foxtail during the early growth stages or when the foxtail is 1 to 4 inches tall. Use two applications with a 14-day interval between applications.
- **Yellow nutsedge:** Two applications are required when the yellow nutsedge plants are 3 to 6 inches tall. Do not apply the second application until 14 days after the first application.
- **Dandelion, plantain, buttercup, speedwell, knotweed, pennywort (dollarweed) and other broadleaf weeds:** Apply when broadleaf weeds are young and actively growing with adequate soil moisture.

FALL:

- **Henbit, chickweed, white clover, ground ivy, lawn burweed, pearlwort, Carolina geranium and other broadleaf weeds:** Apply when broadleaf weeds are young and actively growing with adequate soil moisture for best results.

REFILL CONTAINER

Use this product to refill the empty Roundup® For Lawns$_1$ container by pouring product carefully and directly into the container. DO NOT add water.

HOW TO REFILL

Do not reuse or refill this container. Use this product to refill the empty Roundup® For Lawns$_1$ container by pouring product carefully and directly into the container. DO NOT add water.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE:	Store product in original container in a safe place away from direct sunlight. Keep from freezing. If frozen, allow to thaw and shake well before using.
PESTICIDE DISPOSAL AND CONTAINER HANDLING:	Nonrefillable container. Do not reuse or refill this container unless the directions for use allow a different concentrated or ready-to-use product to be diluted in or poured directly into the container. Reuse or refill this container according to the directions contained in the **HOW TO REFILL** section. **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
IF IN EYES:	• Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-246-7219 for emergency medical treatment advice.	

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff may be hazardous to aquatic organisms in water and to plants adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

† **GUARANTEED RESULTS:** If for any reason you are not satisfied after using this product, simply send us original proof of purchase and we will refund the purchase price.

INSTRUCCIONES DE USO

Utilizar este producto de modo inconsistente con las declaraciones de su etiqueta constituye una violación de las leyes federales.

RESTRICCIONES:



- No aplique este producto de forma que entre en contacto con otras personas o mascotas, ya sea directamente o al ser arrastrado por el viento. No permita que las personas o las mascotas ingresen al área tratada hasta que se seque el rocío.

PRECAUCIONES:

Roundup® For Lawns, es un producto listo para usar (RTU) indicado para los propietarios de casas para el tratamiento "localizado" de grama y malezas de hoja ancha en céspedes. Este producto debe aplicarse a los céspedes con las siguientes especies de grama: azul de Kentucky, raigrás perenne, especies de cañuela o festuca (incluyendo las variedades alta, rubra y roja), Bermuda[1], búfalo y zoysia.

[1] La aplicación en el césped Bermuda puede causar amarillamiento o decoloración temporal, pero se puede esperar una recuperación completa. No aplique el producto a césped Bermuda cuando las temperaturas diurnas superen los 85° F.

Roundup® For Lawns, **no** debe utilizarse en céspedes de las variedades agrostis, San Augustín, bahía, ciempiés o alfombra. **No** lo utilice en dicondra ni en céspedes con tréboles o leguminosas que no desee eliminar, huertos de hortalizas, o plantas frutales u ornamentales (flores, árboles, cubiertas vegetales, setos, macizos de flores o jardines, y arbustos).

CUÁNDO USARLO

Para obtener los mejores resultados, aplique este producto a las malezas durante las primeras etapas de crecimiento activo. Pueda que se necesiten dos o más aplicaciones, a intervalos de 14 días, durante toda la temporada para eliminar las malezas gramíneas. Aplique cuando la temperatura diurna oscile entre los 45° y los 90° F.

EN LA PRIMAVERA Y A PRINCIPIOS DEL VERANO:

- **Para tratar el zacate mano de cangrejo (variedades lisa y sangradera ancha):** Aplique este producto a partir de la germinación del zacate mano de cangrejo hasta la etapa de 6 hojas (2 brotes) o cuando alcance una altura de 3 a 4 pulgadas. Este período comienza a partir del momento en el que observe zacate mano de cangrejo en el césped. Realice dos aplicaciones, dejando un intervalo de 14 días entre una y otra.
- **Para tratar el zacate cola de zorra (variedades verde, gigante y amarilla):** Aplique este producto al zacate cola de zorra durante las primeras etapas de crecimiento o cuando tenga entre 1 y 4 pulgadas de alto. Use dos aplicaciones, dejando un intervalo de 14 días entre una y otra.
- **Para tratar la juncia avellanada:** Es necesario hacer dos aplicaciones cuando las plantas de juncia avellanada tengan entre 3 y 6 pulgadasde alto. No realice la segunda aplicación hasta 14 días después de la primera.
- **Para tratar el diente de león, el lantén, la celidonia menor, la Verónica, el polígono, la centella asiática (acaricoba) y otras malezas de hoja ancha:** Aplique este producto cuando las malezas de hoja ancha sean pequeñas y estén creciendo de manera activa, con un suelo que tenga la humedad adecuada.

EN EL OTOÑO:

- **Para tratar la ortiga mansa, la estalaria, el trébol blanco, la hiedra rastrera, el codillo prieto del césped, la sagina, el geranio silvestre y otras malezas de hoja ancha:** A fin de obtener los mejores resultados, aplique este producto cuando las malezas de hoja ancha sean pequeñas y estén creciendo de manera activa, con un suelo que tenga la humedad adecuada.

RECARGA DE CONTENEDORES

Utilice este producto para rellenar el envase vacío del producto Roundup® For Lawns, vertiendo el producto con cuidado directamente en el contenedor. NO agregue agua.

CÓMO RELLENAR

No reutilice o rellene este envase. Utilice este producto para rellenar el envase vacío del producto Roundup® For Lawns, vertiendo el producto con cuidado directamente en el contenedor. NO agregue agua.

ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua, la comida ni el pienso para animales al almacenar o eliminar el producto.

ALMACENAMIENTO DEL PESTICIDA:	Guarde el producto en su envase original en un lugar seguro y alejado de la luz directa del sol. Proteja el producto del congelamiento. Si se congela, deje que se descongele y agítelo bien antes de usarlo.
ELIMINACIÓN DEL PESTICIDA Y MANEJO DEL ENVASE:	Este envase no se puede rellenar. No vuelva a usar o a rellenar este envase, a menos que las instrucciones de uso permitan que un producto concentrado o listo para usar distinto se pueda echar directamente o diluir en el envase. Este envase se puede reutilizar o rellenar de acuerdo con las instrucciones contenidas en la sección **CÓMO RELLENAR**. **Si está vacío:** Tírelo a la basura o recíclelo, de ser posible. **Si está parcialmente lleno:** Llame a su agencia local de desechos sólidos para obtener instrucciones sobre cómo eliminarlo. Nunca elimine el producto no utilizado en drenajes internos o externos.

AVISOS DE PRECAUCIÓN

RIESGOS PARA LOS SERES HUMANOS Y LOS ANIMALES DOMÉSTICOS

PRECAUCIÓN: Causa irritación moderada en los ojos. Evite el contacto con los ojos o la ropa. Lávese con abundante agua y jabón después de manipular el producto y antes de comer, beber, masticar chicle, consumir tabaco o usar el inodoro.

PRIMEROS AUXILIOS	
SI ENTRA EN LOS OJOS:	• Mantenga los ojos abiertos y enjuáguelos lentamente y con cuidado con agua por 15 a 20 minutos. Retire los lentes de contacto, de haberlos, después de trascurridos los primeros 5 minutos, y continúe enjuagando el ojo. • Llame a un centro de control de envenenamientos para recibir instrucciones de tratamiento.
Tenga el envase o la etiqueta del producto con usted cuando llame al centro de envenenamientos o al médico, o al acudir a recibir tratamiento. También puede llamar al 1-800-246-7219 para obtener información sobre tratamiento médico de emergencia.	

RIESGOS MEDIOAMBIENTALES

Este producto es tóxico para los peces y los invertebrados acuáticos, así como para las plantas acuáticas. No lo aplique directamente al agua, en áreas donde haya agua presente en la superficie ni en zonas intermareales por debajo de la línea de la marea alta. El producto que se escurra o sea arrastrado por el viento puede ser peligroso para las plantas y para los organismos acuáticos de los cuerpos de agua adyacentes a las superficies tratadas. No contamine el agua al deshacerse de la que use para lavar o enjuagar los utensilios. Para proteger el medio ambiente, no permita que el pesticida penetre o se escurra en drenajes pluviales, zanjas de desagüe, cunetas o aguas superficiales. La aplicación de este producto en un clima tranquilo, cuando no se pronostiquen lluvias para las próximas 24 horas, ayudará a asegurar que ni el viento ni la lluvia arrastren o laven el pesticida fuera del área de tratamiento. Enjuagar el equipo usado para la aplicación sobre el área tratada ayudará a evitar escurrimientos hacia cuerpos de agua o sistemas de drenaje.

ADVERTENCIA: En la medida contemplada por las leyes correspondientes, el comprador asume toda la responsabilidad por la seguridad y el uso de este producto si no se siguen las instrucciones.

† **RESULTADOS GARANTIZADOS:** Si por algún motivo usted no queda satisfecho después de usar este producto, simplemente envíenos el comprobante de compra original y le reembolsaremos el precio de compra.

Roundup FOR LAWNS, ®

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
IF IN EYES:	• Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-246-7219 for emergency medical treatment advice.	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE:	Store product in original container in a safe place away from direct sunlight. Keep from freezing. If frozen, allow to thaw and shake well before using.
PESTICIDE DISPOSAL AND CONTAINER HANDLING:	Nonrefillable container. Do not reuse or refill this container unless the directions for use allow a different concentrated or ready-to-use product to be diluted in or poured directly into the container. Reuse or refill this container according to the directions contained in the **HOW TO REFILL** section. **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff may be hazardous to aquatic organisms in water and to plants adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Questions & Comments **Preguntas y comentarios**
Call 1-800-246-7219 or Visit our Website at www.roundup.com

Produced for:
Monsanto Company
Lawn & Garden Products
800 N. Lindbergh Blvd.
St. Louis, MO 63167
© 2020 Monsanto Company

EPA Reg. No. 2217-917-71995
EPA Est. No. 239-IA-3
Roundup and Roundup & Design are registered trademarks of Monsanto Technology LLC.

1005

PPN 10011U1-05
LB15469

FPO

0 70183 50201 1

PRESS TO RESEAL. / PRESIONE PARA VOLVER A CERRAR.