

PROFESSIONAL TURF PRODUCTS

0.48% BARRICADE® ON DG PRO®







PRE-EMERGENT WEED CONTROL

FOR LAWN & LANDSCAPE ORNAMENTALS



Patented Dispersing Granule Technology



Controls over 30 grassy and broadleaf weeds



Small particle formulation provides more particles per square inch

ACTIVE INGREDIENT: Prodiamine* ...0.48% OTHER INGREDIENTS: .99.52% 100.00%

*(CAS No. 29091-21-2)

EPA Reg. No. 9198-214 EPA Est. No. 9198-0H-1<u>M</u>, 9198-0H-2<u>B</u>, 9198-AL-001<u>A</u> Underlined letter is first letter used in run code.

For preemergence control of grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens), lawns and sod nurseries landscape ornamentals
- established perennial & wildflower plantings
- non-crop areas on or surrounding managed rights-of-way for transportation systems
- and utilities (including roadways, roadsides, railways & equipment yards)
- $\bullet \ facilities \ including \ substations, \ tank farms, \ pumping \ stations, \ parking \ \& \ storage \ areas,$ ungrazed fencerows

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BACK PANEL AND BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND FIRST AID

COVERS UP TO 5,800 SQ. FT. NET WT. 18.00 LBS (8.16 KG)



PROFESSIONAL TURF PRODUCTS

0.48% BARRICADE® ON DG PRO®

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

GENERAL INFORMATION

This product is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens), lawns, sod, nurseries, landscape ornamentals, established perennial & wildflower plantings, non-crop areas on or surrounding managed rights-of-way for transportation systems and utilities (including roadways roadsides, railways & equipment yards), facilities including substations, tankfarms, pumping stations, parking & storage areas,

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Do not graze or feed livestock forage cut from areas treated with this product. Do not apply this product through any type of irrigation system unless otherwise instructed by this label. Do not apply aerially. Do not apply to golf course putting greens.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty container into application equipment. Then dispose of empty bag or box in a sanitary landfill or at an incineration facility; or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

APPLICATION DIRECTIONS

Apply uniformly through suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map to the below for approximate crabgrass seed germination dates.

CRABGRASS GERMINATION DATES

Turf Species

MAXIMUM APPLICATION RATE OF THIS PRODUCT PER CALENDAR YEAR BY TURF GRASS SPECIES¹

Lbs. Ai/A Lbs. Product/A Lbs. Product/1,000 sq. ft. Bermudagrass², Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including 312 7.1 turftype), Zoysia Buffalograss, Kentucky Bluegrass, Perennial Ryegrass 4.7 208 Creeping Red Fescue 156 3.5 Creeping Bentgrass 0.65 135

These are the maximum rates per calendar year by species limitations.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 108 lb./acre or 2.4 lb./1,000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

Use higher rates of this product to achieve higher levels weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates Table.

These suggested spreader settings are not intended to replace calibration. Calibrate your spreader before applying product.

18 lbs. treats 5,800 sq. ft. at 3.1 lbs/1,000 sq. ft. rate

18 lbs. treats 3,800 sq. ft. at 4.7 lbs/1,000 sq. ft. rate

18 lbs. treats 5.100 sq. ft. at 3.5 lbs/1.000 sq. ft. rate 18 lbs. treats 2.500 sq. ft. at 7.1 lbs/1.000 sq. ft. rate

10 103. treats 3,100 3q. 1t. at 3.3 103/1,000 3q. 1t. fate		10 103. treats 2,500 5q. rt. at 7.1 103/1,000 5q. rt. rate		
SPREADER MODEL	LOW RATE 3.1 lbs./1,000 sq. ft	MEDIUM RATE 3.5 lbs./1,000 sq. ft	HIGH RATE 4.7 lbs./1,000 sq. ft	HEAVY RATE 7.1 lbs./1,000 sq. ft
The Andersons LCO-1000	K 1/2	L	M 1/2	Р
The Andersons Yard Star 2150	13 1/2	14	15 1/2	18
Agri-Fab Broadcast (1-10)	3 1/2	4	5	7
Agri-Fab Drop (0-10)	3 1/2	4	5	7
Earthway Broadcast (0-30)	13 1/2	14	15 1/2	18
Earthway Drop (0-30)	13 1/2	14	15 1/2	18
Earthway 3400 Hand Spreader (1-3)	1	2	3	-
Scotts Broadcast (2-15)*	4	4 1/2	51/2	7 1/2
Scotts Drop**	6 1/4	6 1/2	7 1/2	9 1/2

*Includes EdgeGuard, Elite, Standard, Basic, LawnPro & Speedy Green 1000 - 3000

**Accugreen 1000-3000 & Classic

DGRR18 1 ROOK

Peel Here

RATES OF APPLICATION This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table.

Product of U.S.A MANUFACTURED BY: The Andersons, Inc. PO Box 119 | Maumee, OH 43537 info@andersonshomeandgarden.com 877-474-8262

QUESTIONS?

1 - After May 30

2 - After May 10 3 - After April 20 4 - After March 20

5 - Jan.1 to March 20

