COVERS 6,000 SQ FT

NET WT/PESO NETO
18.78 lb (8.51 kg)

ACTIVE INGREDIENT:
Metsulfuron-methyl† 0.022%
OTHER INGREDIENTS: 99.978%
TOTAL 100.000%
†CAS #74223-64-6

KEEP OUT OF REACH OF CHILDREN
CAUTION SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN CONSULTE EL PANEL POSTERIOR PARA CONOCER LOS AVISOS DE PRECAUCIÓN ADICIONALES.

32-0-9

DIRECTIONS FOR USE
INSTRUCCIONES DE USO

It is a violation of federal law to use this product in a manner inconsistent with its labeling.
This product applies 0.03 lb active ingredient per acre.
Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.
Con este producto se aplican 13.6 gramos del ingrediente activo por cada 4047 m².

Scotts® QUICK START GUIDE: You're Just Three Simple Steps Away From A Beautiful Scotts® Lawn!
Guía de inicio rápido de Scotts®: ¡Está a sólo tres sencillos pasos de tener un hermoso césped con Scotts®!

FILL & SET SPREADER
Llene y ajuste el esparcidor

Scotts® Broadcast / Rotary Spreaders **3¼**

Scotts® Drop Spreaders **5½**

This package applies 3.13 lb of product per 1000 sq ft.

Con este paquete, se aplican 1.4 kg de producto por cada 93 m².

APPLY
Aplique

- For use on St. Augustinegrass (including Floratam), bermudagrass, centipedegrass, zoysiagrass, seashore paspalum and buffalograss.
- May be used on tall fescue, Kentucky bluegrass and fine fescues if temporary discoloration can be tolerated.
- Apply when weeds are actively growing.
- May be applied to a wet or dry lawn.
- For best results apply to grass that is wet from dew or watering.
- Mow tall grass a day or two before application.
- If rain does not occur within 2-3 days after application, water in lightly for activation.
- For even application walk at a steady pace slightly overlapping each pass.
- Scotts® Drop Spreaders: Align hopper arrow with previous wheel track.
- Scotts® Broadcast/Rotary Spreaders: Line up each pass approximately 5 ft. from wheel track.

- Para su utilización en céspedes San Agustín (incluyendo la variedad Floratam), Bermuda, ciempiés, zoysia, del mar o chepica blanca, y búfalo.
- Puede usarse en festuca o cañuela alta, hierba azul de Kentucky y festuca o cañuela rubra, si una decoloración temporal es tolerable.
- Aplique cuando las malezas estén creciendo de manera activa.
- Puede aplicarse al césped tanto húmedo como seco.
- Para obtener los mejores resultados, aplique el producto al césped humedecido con agua de rocío o de riego.
- Corte el césped alto uno o dos días antes de la aplicación.
- En caso de que no llueva en dos a tres días después de la aplicación, riegue ligeramente para activar el producto.
- Para que la aplicación sea uniforme, camine a un paso constante, traslapando ligeramente cada pasada.
- Esparcidores por goteo Scotts®: Alinee la flecha de la tolva con la huella dejada previamente por la rueda.
- Esparcidores por voleo/giratorios Scotts®: Alinee cada pasada a aproximadamente 1.5 m de la huella dejada por la rueda.

CLEAN-UP
Limpie

Sweep product from hard surfaces onto the lawn.

Barra el producto derramado sobre superficies duras hacia el césped.

YOU'RE DONE! WHAT'S NEXT?
¡Listo! ¿Y ahora qué sigue?

Do not allow people or pets to enter treatment areas until the application is complete and the grass is dry.

No deje que las personas o las mascotas ingresen a las áreas de tratamiento hasta que termine de aplicar y el césped esté seco.

Questions? Call
1-800-543-TURF
¿Preguntas? Llame al 1-800-543-TURF.

www.scotts.com

RESTRICTIONS
- Do not use on bahiagrass, ryegrass (annual and perennial) and blends that contain ryegrass.
- Do not apply this product in a way that will contact any person either directly or through drift. Keep people and pets out of area during application.
- Do not apply on or under the branch spread area of trees, shrubs, bedding plants, flowers or garden plants.
- Do not apply more than two times per year.

SEEDING/SPRIGGING
- May be applied after sprigging/sodding a new St. Augustinegrass, bermudagrass, centipedegrass or zoysiagrass lawn to prevent competition from germinating weeds. Water the new sprigs, then make the application of this product.
- Do not apply to newly seeded lawns until after the new grass has been mowed at least three times.
- Do not plant sprigs/sod for one month, or sow new grass seed for two months following application of this product.

RESTRICCIONES
- No aplique en céspedes bahía o raigrás (ya sea anual o perenne), ni en mezclas que contengan raigrás.
- No aplique este producto de forma que entre en contacto con otras personas, ya sea directamente o al ser arrastrado por el viento. Durante la aplicación, mantenga alejadas del área a las personas y las mascotas.
- No aplique sobre ni debajo del área abarcada por las ramas de árboles, arbustos, plantas de macizo, flores o plantas de jardín.
- No aplique más de dos veces por año.

SEMBRADO DE SEMILLAS/RETOÑOS
- Puede aplicarse después de plantar retoños o cubrir con césped San Agustín, Bermuda, ciempiés o zoysia un nuevo prado, para evitar que compita con malezas en germinación. Riegue los nuevos retoños y luego aplique este producto.
- No aplique a prados recién sembrados sino hasta después de por lo menos el tercer corte de césped.
- No plante retoños ni tepes por un mes, ni siembre semillas de césped por dos meses, después de la aplicación de este producto.

WEEDS CONTROLLED

Controls
Annual sowthistle
Aster
Black medic
Broadleaf plantain
Buckhorn plantain
California burclover
Carolina geranium
Catsear
Chicory
Common chickweed
Common groundsel
Common mallow
Common purslane
Common yarrow
Corn speedwell
Creeping woodsorrel
Crown vetch
Cudweed
Dandelion
Dollarweed (pennywort)
Eclipta
English daisy
Evening primrose
Florida betony
Florida pusley
Ground ivy
Hairy bittercress
Henbit
Hop clover
Horseweed
Khakiweed
Lambsquarters
Lawn burweed (spurweed)
Mousear chickweed
Mustard
Parsley-piert
Pensacola bahiagrass
Perennial ryegrass
Pigweed
Pineapple-weed
Prostrate knotweed
Puncturevine
Purple deadnettle
Red clover
Shepherd's-purse
Spurge
White clover
Wild carrot
Wild garlic
Wild violet
Yellow woodsorrel *(Oxalis)*

Suppresses
Argentine bahiagrass
Canada thistle
Field bindweed
Virginia buttonweed

2-Month Control
Broadleaf plantain
Buckhorn plantain
California burclover
Dandelion
Dollarweed (pennywort)
White clover

TIPS FOR LAWNCARE SUCCESS

Feed Regularly
Regular feedings provide the nutrients your lawn needs to look its best and withstand the stresses of weeds, heat, drought & family activity.

Benefits of Multiple Feedings

1 ANNUAL FEEDING *2-3 ANNUAL FEEDINGS* *4 ANNUAL FEEDINGS*

Mow and Leave the Grass Clippings on Your Lawn
Mowing high allows the grass to develop a deep root system & helps your lawn use water more effectively. Grass clippings recycle the plant nutrients back into the soil.

Conserve Water
Well-fed lawns are more efficient at using available water. Feeding with Scotts® lawn food helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn against heat, drought and other stresses. Remember your lawn will begin to wilt when water is needed. As much as possible take advantage of nature's sprinkler and rely on the rain to water your lawn.

SCOTTS® RECOMMENDED FEEDING ROUTINE

EARLY SPRING
An early spring feeding helps grass get going after its initial green up. You should feed around the first time you mow your grass.

LATE SPRING
Feeding now prepares grass for summer by making more blades on top and building deep roots below.

SUMMER
A summer feeding provides grass the nutrients they need to thrive during the summer months.

EARLY FALL
Feeding now helps grass store nutrients to deal with the stress of the winter months.

Florida Applications: 4 annual feedings with the exception of centipedegrass (2 feedings in the north and 3 feedings in the central and south).

North Florida is north of Ocala. Central Florida is defined as south of Ocala to a line extending from Vero Beach to Tampa. South Florida includes the remaining southern portion of the state.

Scotts & Your Environment

We care. Scotts® products are designed with care to grow thicker, greener lawns. **A Green Lawn Is a Good Lawn -** A healthy lawn can clean the air, produce oxygen and prevent runoff and soil erosion. **You can make a difference -** Help keep our water resources clean. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk or street back onto your lawn.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals
CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment. Call 1-800-543-8873.

ENVIRONMENTAL HAZARDS: To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treated area. Sweeping any product that lands on a driveway, sidewalk, or street, back into the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

STORAGE AND DISPOSAL
PESTICIDE STORAGE: Store in a dry place. Reseal opened bag by folding top down and securing. Store in original container in a secure area away from children and pets.
PESTICIDE DISPOSAL and CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.
If Empty: Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

AVISOS DE PRECAUCIÓN Riesgos para humanos y animales domésticos
PRECAUCIÓN: Causa irritación moderada en los ojos. Evite el contacto con los ojos o la ropa. Lávese con abundante agua y jabón después de manipular el producto y antes de comer, beber, mascar chicle, consumir tabaco o usar instalaciones sanitarias.

PRIMEROS AUXILIOS
SI ENTRA EN LOS OJOS: Mantenga el ojo abierto y enjuague con agua despacio y con cuidado por 15 a 20 minutos. Si tuviera lentes de contacto, quítelos después de lavarse con agua los primeros 5 minutos y luego continúe enjuagando el ojo. Llame al centro de control de envenenamientos o al médico para obtener recomendaciones de tratamiento. Tenga el envase del producto o la etiqueta con usted cuando llame a un centro de control de envenenamientos o al médico, o al acudir a recibir tratamiento. Llame al 1-800-543-8873.

RIESGOS AMBIENTALES: Para proteger el medio ambiente, no permita que el pesticida penetre o se escurra en drenajes pluviales, zanjas de desagüe, cunetas o aguas superficiales. La aplicación de este producto en un clima tranquilo, cuando no haya un pronóstico de lluvia para las próximas 24 horas, asegurará que el viento o la lluvia no soplen o laven el pesticida fuera del área de tratamiento. Barra el producto que caiga sobre la entrada de automóviles, la acera o la calle hacia el área tratada del césped para evitar que se escurra a cuerpos de agua o sistemas de drenaje.

ALMACENAMIENTO Y ELIMINACIÓN
ALMACENAMIENTO DEL PESTICIDA: Guardar en un lugar seco. Cierre la bolsa abierta doblando la parte superior y asegurándola. Almacene el producto en su envase original, en un lugar seguro y alejado de los niños y las mascotas.
ELIMINACIÓN y MANEJO DEL ENVASE DEL PESTICIDA: Este envase no se puede rellenar. No vuelva a usar ni a llenar este envase. **Si está vacío:** Tírelo a la basura o, si está disponible, ofrézcalo para reciclaje. **Si está parcialmente lleno:** Llame a su agencia local de desechos sólidos para obtener instrucciones sobre cómo eliminarlo. Nunca elimine el producto no utilizado en drenajes internos o externos.

Scotts®Turf Builder® Bonus® S Southern Weed & Feed 32-0-9 F643

GUARANTEED ANALYSIS
Total Nitrogen (N) 32%
17.6% Urea Nitrogen
13.2% Other Water Soluble Nitrogen*
1.2% Water Insoluble Nitrogen*
Soluble Potash (K2O) 9%

Derived from Methyleneureas, Urea, Potassium Chloride.
*Contains 10.8% slowly available Nitrogen from Methyleneureas.

EPA Reg. No. 538-323, EPA Est. No. 538-OH-1[S], 35512-FL-2[H], 85653-OH-1[R]
Superscript used is first letter of run code on end of container.

QUESTIONS / INFORMATION
Preguntas / Información

SMG Listed NYSE
Sold and Guaranteed By:
The Scotts Company
14111 Scottslawn Rd.
Marysville, OH 43041

Questions? Call
1-800-543-8873

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-sc.com

La información relacionada con los contenidos y los niveles de metales de este producto está disponible en Internet en http://www.regulatory-info-sc.com

©2014 The Scotts Company LLC
World rights reserved.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

AVISO: En la medida contemplada por las leyes aplicables, el comprador asume todos los riesgos de uso, almacenamiento o manejo de este producto si no se siguen las instrucciones.

SCOTTS NO-QUIBBLE GUARANTEE®
If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund promptly.

Si por alguna razón no está satisfecho luego de utilizar este producto, usted, el consumidor, tiene derecho a la devolución de su dinero. Sólo envíenos el comprobante de compra y le enviaremos un reembolso por correo inmediatamente.

PROOF OF PURCHASE UPC 0-32247-33135-7
33135-2 VID-BB BG4111

FPO

0 32247 33135 7

SCOTTS® TURF BUILDER® BONUS® S SOUTHERN WEED & FEED
GM 33135-2