



quick FACTS

- KILLS BROADLEAF WEEDS INCLUDING DOLLARWEED, DANDELION, CLOVER, CHICKWEED & HENBIT**
- PREVENTS GRASSY AND BROADLEAF WEEDS SUCH AS CRABGRASS, ANNUAL BLUEGRASS, COMMON DANDELION, GOOSEGRASS & SPURGE**
- FEEDS & STRENGTHENS THE LAWN. CONTAINS SLOW RELEASE FERTILIZER FOR LONG LASTING FEEDING.**



WHEN TO APPLY

Apply anytime in early spring or fall.

WHERE TO USE

Use only on established lawns of the turf types listed below:

Bermudagrass, Centipede grass, all varieties of St. Augustine grass including Floratam, and Zoysia
DO NOT use on Ormond variety of Bermudagrass, Fescue, Bentgrass, Carpetgrass, Kikuyugrass, Ryegrass or Dichondra lawns

COVERAGE This bag treats 5,000 sq. ft.



Questions or comments, call toll-free (877) 229-3724
www.BayerAdvanced.com

For the full Spanish translation go to www.BayerAdvanced.com



Tired of killing weeds in your lawn – even after you've put down your traditional weed & feed granules? Just one application of Bayer Advanced 3-in-1 Weed & Feed for Southern Lawns kills lawn weeds and prevents new ones, including Crabgrass, for up to 6 months. Plus it greens, feeds and strengthens your lawn for a thick, healthy turf. It delivers season long weed control that is simple and easy to use and it will not harm your lawn when used as directed. No need for frequent re-application. Plus, when applied in early spring, it prevents Crabgrass. And when re-applied late summer, prevents Annual Bluegrass. Up to 6 months of weed control – plus feeding – has never been easier.

***Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE

- Use this product only on established Southern lawns – new lawns must be at least 3 months old if sodded, 16 months old if sprigged & 16 months old if seeded.
- Do not overlap treated areas or make repeat applications within 90 days as over application could damage the lawn.
- Apply only to actively growing turf.
- Do not apply to recently aerated or verticulated turf until it has recovered.
- Apply when the air is calm to avoid drift onto desirable plants.

HOW TO USE

- Use rotary or drop type spreader to apply granules evenly over the lawn. To be sure you cover lawn areas uniformly, first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules.
- Sweep, brush, or blow any accidentally applied material off driveways, sidewalks etc. back onto the lawn.
- Wait 24 hours after applying then activate residual barrier by lightly watering in.
- Do not make more than two applications per site in any 12 month period.
- Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.

FOR BEST RESULTS

WHEN TO APPLY

- Apply when - weeds are small and actively growing - temperatures are between 50–90°F.
- May be applied to dry or (for best results) wet lawns.
- Do not apply while turfgrass is emerging from dormancy.
- Do not apply to Centipede grass after October 1.
- To prevent Crabgrass and Goosegrass apply early spring before the temperatures are regularly in the lower 80° F range.
- To prevent Annual Bluegrass apply August - December.
- Lawns may be seeded 12 months, sodded 6 months, and sprigged 12 months after applying this product.

MOWING

- If the lawn needs mowing, do it 1 to 2 days before (for best results) or 1 day after applying this product.
- Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

WATERING

- If lawn is dry, water 1 to 2 days before using this product.
- Do not water for at least 24 hours after application or apply if rain is expected within 24 hours.

HOW IT WORKS

This product contains a proven blend of weed killers to kill existing weeds plus a residual herbicide to keep them from coming back for up to 6 months. This product starts working immediately to stop weed growth until weeds are dead within weeks. Plus, it feeds your lawn for quick greening and long lasting feeding.




Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

SPREADER SETTINGS

If your spreader is not listed, please call our Consumer Service Hotline toll-free (877) 229-3724 for help.

- Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.
- Tip: 1,000 sq ft = 20 ft x 50 ft
- Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.



SETTING TO DELIVER 2.5lb/1000 sq. ft.

Broadcast Spreader	SETTING
Earthway Ev-N-Spread	10
Scotts Basic	3
Scotts Deluxe Edgeguard	3
Scotts Mini	3
Scotts Standard	8
Sta-Green	3
Drop Spreader	
Scotts Accugreen 1000	2.5
Scotts Accugreen 3000	2

KILLS

Beggarweed, Creeping Beggarweed, Dwarf Betony, Florida Burdock
Buttercup, Hairy Buttonweed, Virginia* Clover, Hop Clover, Red Cocklebur, Common Coffeeweed, (Common Chickory) Daisy, English (Lawn Daisy) Dandelion, False

Dollarweed (Pennywort) Goldenrod Hawkweed Heallall Ivy, Ground (Creeping Charlie) Knotweed Kyllinga, Green* Lespedeza, Common Mallow, Roundleaf Matchweed Nutsedge, Purple* Nutsedge, Yellow* Parsley-piert

Peppergrass Pepperweed, Virginia Plantain, Broadleaf Smartweed Sorrel, Sheep Speedwell, Common Spotted Catsear Spurweed Thistle Vetch, Common Wild Violet Woodsorrel, Creeping

* Suppression

PREVENTS

Ageratum, Tropic Barley, Mouse Barnyardgrass Crabgrass, Smooth Cudweed, Linear Leaf (Purple Cudweed) Dandelion, Cat's Ear Doveweed Evening Primrose, Common Evening Primrose, Cutleaf Flatsedge, Rice Fleabane, Hairy

Crabgrass, Large (Hairy Crabgrass) Crabgrass, Smooth Cudweed, Linear Leaf (Purple Cudweed) Dandelion, Cat's Ear Doveweed Evening Primrose, Common Evening Primrose, Cutleaf Flatsedge, Rice Fleabane, Hairy

Foxtail, Giant Foxtail, Green Foxtail, Yellow Filaree, Redstem Parthenium Groundsel, Common Goosegrass Kochia Kyllinga, Fragrant Lovegrass, Tufted Mallow, Little Maretail Mulberry Weed

Mustard, Black Mustard, Wild Nightshade, American Black Nightshade, Hairy Parthenium Phyllanthus, Longstalked Pigweed, Prostrate Pigweed, Redroot Spindash, Palesseed Puncturevine Redmaids Rocket, London Ryegrass, Italian

Ryegrass, Perennial Sandbur Sedge, Annual Sesbania, Hemp Sida, Prickly (Teaweed) Southern Brassbuttons Sowthistle, Annual Spindash, Corn Spurge, Prostrate Sunflower, Common Swinecress Tassel Flower

Velvetleaf Vetch, Purple Wild Carrot Willowherb Woodsorrel (Oxalis)

Burnweed, American Chickweed, Common Chickweed, Mouseear Clover, White Dandelion, Common Dock, Curly Eclipta Henbit Knotweed, Prostrate Kyllinga, Annual* Lambsquarters Medic, Black Plantain, Buckhorn

* Suppression

ENVIRONMENTAL HAZARDS

- This product is toxic to fish, aquatic invertebrates, and plants. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean water mark. Do not contaminate water when disposing of rinsate or washwater. This product may impact water through spray drift or runoff. Follow directions for use to avoid spray drift and runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams and springs will reduce the potential of this product entering water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.
- **Ground Water Advisory:** This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.
- **Surface Water Advisory:** This pesticide may impact water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

PELIGROS AMBIENTALES • Este producto es tóxico para los peces, los invertebrados acuáticos y las plantas. No aplicar directamente al agua o en áreas donde haya aguas superficiales o áreas entre mareas por debajo de la marca de agua. No contaminar el agua al eliminar el agua de lavado o enjuague. Este producto puede afectar el agua a través del rocío o derrames. Seguir las instrucciones de uso para evitar rocíos y derrames. Un buen mantenimiento de la zona de contención de vegetación entre las áreas en las que se aplica este producto y la diferenciación del agua de superficie, incluidos estanques, arroyos y manantiales, reduce el potencial de penetración en el agua a causa de precipitación-derrame. El derrame de este producto se reducirá al evitar las aplicaciones cuando se pronostica lluvia dentro de las 48 horas. • **Recomendación sobre las napas subterráneas:** Este pesticida contiene propiedades y características relacionadas con sustancias químicas detectadas en aguas subterráneas. Este producto químico puede filtrarse en las napas subterráneas si se utiliza en áreas donde los suelos son permeables, particularmente donde el nivel freático es poco profundo. • **Recomendación sobre el agua de superficie:** Este pesticida puede impactar sobre la calidad del agua a causa de derrame de agua pluvial. Esto se cumple especialmente en suelos con drenaje deficiente y suelos con napas subterráneas poco profundas. Este producto se clasifica como con alto potencial para llegar al agua de superficie a través de derrames durante varios meses o más después de la aplicación.

FIRST AID



NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information. Active Ingredients: 0.088% Dicamba, (CAS n.°1918-00-9), 0.041% Penoxsulam (CAS n.°219714-96-2) and 0.027% Indaziflam (CAS n.°950782-86-2).

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 – 20 minutes.
- Call a poison control center or doctor for further treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for further treatment advice.

PRIMEROS AUXILIOS EN CASO DE INGESTIÓN

• Llamar inmediatamente a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. • Hacer que la persona beba tragos de agua, si puede tragar. • No inducir el vómito salvo que así lo indique el médico o el centro de control de envenenamiento. • No administrar nada por vía oral a una persona inconsciente. **EN CASO DE CONTACTO CON LA PIEL O LA ROPA** • Quitar la ropa contaminada. • Enjuagar la piel inmediatamente con abundante agua durante 15 a 20 minutos. • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. **EN CASO DE CONTACTO CON LOS OJOS** • Mantener abiertos los ojos y enjuagar de forma lenta y suave con agua durante entre 15 y 20 minutos. • Retirar los lentes de contacto, si los hubiera, después de los primeros 5 minutos y luego continuar enjuagando los ojos. • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. **NOTA:** Tener la etiqueta del producto a mano al llamar a un centro de control de envenenamiento. Si busca tratamiento en una sala de emergencias o en el consultorio de un médico, lleve esta etiqueta del producto para mostrársela al personal médico. También puede llamar al número gratuito 1-877-229-3763 para solicitar información médica de emergencia. Ingredientes activos: 0.088% de dicamba, (CAS n.°1918-00-9), 0.041% de penoxsulam (CAS n.°219714-96-2) y 0.027% de indaziflam (CAS n.°950782-86-2).

STORAGE & DISPOSAL



Do not contaminate water, food, or feed by storage and disposal


PESTICIDE STORAGE Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
PESTICIDE DISPOSAL & CONTAINER HANDLING If Empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

ALMACENAMIENTO Y ELIMINACIÓN

No contaminar el agua o los alimentos al almacenar o eliminar

ALMACENAMIENTO DE PESTICIDAS Almacenar en el contenedor original en un lugar fresco y seco, fuera del alcance de los niños, preferentemente en un gabinete bajo llave. Proteger del congelamiento.

ELIMINACIÓN DE LOS PESTICIDAS Y MANIPULACIÓN DE ENVASES Si está vacío: Envase no rellenable. No rellenar ni volver a utilizar este envase. Colocar el envase vacío en la basura o entregarlo para reciclar si fuera posible. Si está parcialmente lleno: Llamar a la agencia local de residuos sólidos o al número gratuito 1-877-229-3724 para obtener indicaciones para la eliminación. Nunca colocar el producto sin utilizar debajo de algún desagüe interno o externo.



EPA Reg. No. 72155-112

EPA Est. No. indicated by 3rd and 4th digits of the batch number on this package.
(26) = 82757-FL-1 (57) = 071106-GA-003

Bayer Advanced 3-in-1 Weed & Feed for Southern Lawns 35-0-3

GUARANTEED ANALYSIS

Total Nitrogen (N) 35.0%
35% Urea Nitrogen*
Soluble Potash (K₂O) 3.0%
Sulfur (S) Total 4.6%
3.5% Free Sulfur (S)
1.1% Combined Sulfur (S) **F1514**

Derived From: Urea, Polymer Coated Sulfur Coated Urea, & Potassium Sulfate

*Contains 17.5% slowly available Nitrogen from Polymer Coated Sulfur Coated Urea

Information regarding the contents and levels of metals in this product are available on the Internet at <http://www.aapico.org/metals.htm>.

Check with your county or city government to determine if there are local regulations for fertilizer use.

Produced for: Bayer Advanced, A Division of Bayer CropScience LP, P.O. Box 12014, 2 T.W. Alexander Dr., Research Triangle Park, NC 27709

Bayer (reg'd), the Bayer Cross (reg'd) and Bayer Advanced™ are trademarks of Bayer

US449562a 150421A1T

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.